

# ADDITIVE SCHWARZ PRECONDITIONING FOR P-VERSION TRIANGULAR AND TETRAHEDRAL FINITE ELEMENTS

JOACHIM SCHÖBERL, JENS M. MELENK, CLEMENS G. A. PECHSTEIN,  
AND SABINE C. ZAGLMAYR

ABSTRACT. This paper analyzes two-level Schwarz methods for matrices arising from the  $p$ -version finite element method on triangular and tetrahedral meshes. The coarse level consists of the lowest order finite element space. On the fine level, we investigate several decompositions with large or small overlap leading to optimal or close to optimal condition numbers. The analysis is confirmed by numerical experiments for a model problem.

## 1. INTRODUCTION

High order finite element methods can lead to very high accuracy and are thus attracting increasing attention in many fields of computational science and engineering. The monographs [SB91, BS94, Sch98, KS99, SDR04] give a broad overview of theoretical and practical aspects of high order methods.

As the problem size increases (due to small mesh-size  $h$  and high polynomial order  $p$ ), the solution of the arising linear system of equations becomes more and more the time-dominating part. Here, iterative solvers can reduce the total simulation time. We consider preconditioners based on domain decomposition methods [DW90, GO95, SBG96, TW04, Qua99]. The concept is to consider each high order element as an individual sub-domain. Such methods were studied in [Man90, BCM91, Pav94, Ain96a, Ain96b, Cas97, Bic97, GC98, SC01, Mel02, EM04]. We assume that the local problems can be solved directly. On tensor product elements, one can apply optimal preconditioners for the local sub-problems as in [KJ99, BSS04, BS04].

In the current work, we study overlapping Schwarz preconditioners with large or small overlap. The condition numbers are bounded uniformly in the mesh size  $h$  and the polynomial order  $p$ . To our knowledge, this is a new result for tetrahedral meshes. We construct explicitly the decomposition of a global function into a coarse grid part and local contributions associated with the vertices, edges,

---

1991 *Mathematics Subject Classification.* 65N30.

*Key words and phrases.* High Order Finite Element Method, Preconditioning.

The first and the last author acknowledge their support by the Austrian Science Foundation FWF within project grant Start Y-192, “hpFEM : Fast Solvers and Adaptivity”.

faces, and elements of the mesh. The idea of the construction was presented in [SMPZ05].

The rest of the paper is organized as follows: In Section 2 we state the problem and formulate the main results. We prove the 2D case in Section 3 and extend the proof for 3D in Section 4. Finally, in Section 5 we give numerical results for several versions of the analyzed preconditioners.

## 2. DEFINITIONS AND MAIN RESULT

We consider the Poisson equation on the polyhedral domain  $\Omega$  with homogeneous Dirichlet boundary conditions on  $\Gamma_D \subset \partial\Omega$ , and Neumann boundary conditions on the remaining part  $\Gamma_N$ . With the sub-space  $V := \{v \in H^1(\Omega) : v = 0 \text{ on } \Gamma_D\}$ , the bilinear-form  $A(\cdot, \cdot) : V \times V \rightarrow \mathbb{R}$  and the linear-form  $f(\cdot) : V \rightarrow \mathbb{R}$  defined as

$$A(u, v) = \int_{\Omega} \nabla u \cdot \nabla v \, dx \quad f(v) = \int_{\Omega} f v \, dx,$$

the weak formulation reads

$$(1) \quad \text{find } u \in V \text{ such that} \quad A(u, v) = f(v) \quad \forall v \in V.$$

We assume that the domain  $\Omega$  is sub-divided into straight-sided triangular or tetrahedral elements. In general, constants in the estimates depend on the shape of the elements, but they do not depend on the local mesh-size. We define

$$\begin{aligned} \text{the set of vertices} & \quad \mathcal{V} = \{V\}, \\ \text{the set of edges} & \quad \mathcal{E} = \{E\}, \\ \text{the set of faces (3D only)} & \quad \mathcal{F} = \{F\}, \\ \text{the set of elements} & \quad \mathcal{T} = \{T\}. \end{aligned}$$

We define the sets  $\mathcal{V}_f, \mathcal{E}_f, \mathcal{F}_f$  of *free* vertices, edges, and faces not completely contained in the Dirichlet boundary. The high order finite element space is

$$V_p = \{v \in V : v|_T \in P^p \, \forall T \in \mathcal{T}\},$$

where  $P^p$  is the space of polynomials up to total order  $p$ . As usual, we choose a basis consisting of lowest order affine-linear functions associated with the vertices, and of edge-based, face-based, and cell-based bubble functions. The Galerkin projection onto  $V_p$  leads to a large system of linear equations, which shall be solved with the preconditioned conjugate gradient iteration.

This paper is concerned with the analysis of additive Schwarz preconditioning. The basic method is defined by the following space splitting. In Section 5 we will consider several cheaper versions resulting from our analysis. The coarse sub-space is the global lowest order space

$$V_0 := \{v \in V : v|_T \in P^1 \, \forall T \in \mathcal{T}\}.$$

For each inner vertex we define the vertex patch

$$\omega_V = \bigcup_{T \in \mathcal{T}: V \in T} T$$

and the vertex sub-space

$$V_V = \{v \in V_p : v = 0 \text{ in } \Omega \setminus \omega_V\}.$$

For vertices  $V$  not on the Neumann boundary, this definition coincides to  $V_p \cap H_0^1(\omega_V)$ . The additive Schwarz preconditioning operator is  $C^{-1} : V_p^* \rightarrow V_p$  defined by

$$C^{-1}d = w_0 + \sum_{V \in \mathcal{V}} w_V$$

with  $w_0 \in V_0$  such that

$$A(w_0, v) = \langle d, v \rangle \quad \forall v \in V_0,$$

and  $w_V \in V_V$  defined such that

$$A(w_V, v) = \langle d, v \rangle \quad \forall v \in V_V.$$

This method is very simple to implement for the  $p$ -version method using a hierarchical basis. The low-order block requires the inversion of the sub-matrix according to the vertex basis functions. The high order blocks are block-Jacobi steps, where the blocks contain all vertex, edge, face, and cell unknowns associated with mesh entities containing the vertex  $V$ .

The rate of convergence of the cg iteration can be bounded by means of the spectral bounds for the quadratic forms associated with the system matrix and the preconditioning matrix. The main result of this paper is to prove optimal results for the spectral bounds:

**Theorem 1.** *The constants  $\lambda_1$  and  $\lambda_2$  of the spectral bounds*

$$\lambda_1 \langle Cu, u \rangle \leq A(u, u) \leq \lambda_2 \langle Cu, u \rangle \quad \forall u \in V_p$$

*are independent of the mesh-size  $h$  and the polynomial order  $p$ .*

The proof is based on the additive Schwarz theory, which allows to express the  $C$ -form by means of the space decomposition:

$$\langle Cu, u \rangle = \inf_{\substack{u = u_0 + \sum_V u_V \\ u_0 \in V_0, u_V \in V_V}} \|u_0\|_A^2 + \sum \|u_V\|_A^2.$$

The constant  $\lambda_2$  follows immediately from a finite number of overlapping subspaces. In the core part of this paper, we construct an explicit and stable decomposition of  $u$  into sub-space functions. Section 3 introduces the decomposition for the case of triangles, in Section 4 we prove the results for tetrahedra.

### 3. SUB-SPACE SPLITTING FOR TRIANGLES

In this section, we give the proof of Theorem 1 for triangles. The case of tetrahedra is postponed to Section 4.

The strategy of the proof is the following: First, we subtract a coarse grid function to eliminate the  $h$ -dependency. By stepwise elimination, the remaining function is then split into sums of vertex-based, edge-based and inner functions. For each partial sum, we give the stability estimate. This stronger result contains Theorem 1, since we can choose corresponding vertices for the edge and inner contributions (see also Section 5).

**3.1. Coarse grid contribution.** In the first step, we subtract a coarse grid function:

**Lemma 2.** *For any  $u \in V_p$  there exists a decomposition*

$$(2) \quad u = u_0 + u_1$$

such that  $u_0 \in V_0$  and

$$\|u_0\|_A^2 + \|\nabla u_1\|_{L_2}^2 + \|h^{-1}u_1\|_{L_2}^2 \preceq \|u\|_A^2.$$

*Proof.* We choose  $u_0 = \Pi_h u$ , where  $\Pi_h$  is the Clément-operator [Cle75]. The norm bounds are exactly the continuity and approximation properties of this operator.  $\square$

From now on,  $u_1$  denotes the second term in the decomposition (2).

**3.2. Vertex contributions.** In the second step, we subtract functions  $u_V$  to eliminate vertex values. Since vertex interpolation is not bounded in  $H^1$ , we cannot use it. Thus, we construct a new averaging operator mapping into a larger space.

In the following, let  $V$  be a vertex not on the Dirichlet boundary  $\Gamma_D$ , and let  $\varphi_V$  be the piece-wise linear basis function associated with this vertex. Furthermore, for  $s \in [0, 1]$  we define the level sets

$$\gamma_V(s) := \{y \in \omega_V : \varphi_V(y) = s\},$$

and write  $\gamma_V(x) := \gamma_V(\varphi_V(x))$  for  $x \in \omega_V$ . For internal vertices  $V$ , the level set  $\gamma_V(0)$  coincides with the boundary  $\partial\omega_V$  (cf. Figure 1). The space of functions being constant on these sets reads



$$S_V := \{w \in L_2(\omega_V) : w|_{\gamma_V(s)} = \text{const}, s \in [0, 1] \text{ a.e.}\};$$

its finite dimensional counterpart is

$$S_{V,p} := S_V \cap V_p = \text{span}\{1, \varphi_V, \dots, \varphi_V^p\}.$$

We introduce the *spider averaging operator*

$$(\Pi^V v)(x) := \frac{1}{|\gamma_V(x)|} \int_{\gamma_V(x)} v(y) dy, \quad \text{for } v \in L_2(\omega_V).$$

FIGURE 1. The level sets  $\gamma_V(x)$ FIGURE 2. Construction of  $\Pi_0^V$ 

To satisfy homogeneous boundary conditions, we add a correction term as follows (see Figure 2)

$$(\Pi_0^V v)(x) := (\Pi^V v)(x) - (\Pi^V v)|_{\gamma_V(0)}(1 - \varphi_V(x)).$$

**Lemma 3.** *The averaging operators fulfill the following algebraic properties*

(i)

$$\Pi^V V_p = S_{V,p},$$

(ii)

$$\Pi_0^V V_p = S_{V,p} \cap V_V,$$

(iii) *if  $u$  is continuous at  $V$ , then*

$$(\Pi^V u)(V) = \Pi_0^V u(V) = u(V).$$

The proof follows immediately from the definitions.

We denote the distance to the vertex  $V$ , and the minimal distance to any vertex in  $\mathcal{V}$  by

$$r_V(x) := |x - V| \quad \text{and} \quad r_{\mathcal{V}}(x) := \min_{V \in \mathcal{V}} r_V(x).$$

**Lemma 4.** *The averaging operators satisfy the following norm estimates*

(i)

$$\|\Pi^V u\|_{L_2(\omega_V)} \preceq \|u\|_{L_2(\omega_V)}$$

(ii)

$$\|\nabla \Pi^V u\|_{L_2(\omega_V)} \preceq \|\nabla u\|_{L_2(\omega_V)}$$

(iii)

$$\|r_V^{-1}\{u - \Pi^V u\}\|_{L_2(\omega_V)} \preceq \|\nabla u\|_{L_2(\omega_V)}$$

(iv)

$$\|\nabla\{\varphi_V u - \Pi_0^V u\}\|_{L_2(\omega_V)} \preceq \|\nabla u\|_{L_2(\omega_V)}$$

(v)

$$\|r_{\mathcal{V}}^{-1}\{\varphi_V u - \Pi_0^V u\}\|_{L_2(\omega_V)} \preceq \|\nabla u\|_{L_2(\omega_V)}$$

*Proof.* We parameterize the patch  $\omega_V$  by

$$F_V : \gamma_V(0) \times [0, 1] \rightarrow \omega_V : (y, s) \mapsto y + s(V - y).$$

Splitting the patch into elements, and applying element-wise transformation rules, one proves

$$\int_{\omega_V} |f(x)| dx \simeq h_V \int_0^1 \int_{\gamma_V(0)} |f(F_V(y, s))| (1 - s) dy ds,$$

where  $h_V := \text{diam}\{\omega_V\}$ .

(i) Using  $\gamma_V(F_V(y, s)) = \gamma_V(s)$  together with standard inequalities we derive

$$\begin{aligned} \|\Pi^V u\|_{L_2(\omega_V)}^2 &\simeq h_V \int_0^1 \int_{\gamma_V(0)} |(\Pi^V u)(F_V(y, s))|^2 (1 - s) dy ds \\ &= h_V \int_0^1 \int_{\gamma_V(0)} \left| \frac{1}{|\gamma_V(s)|} \int_{\gamma_V(s)} u(x) dx \right|^2 (1 - s) dy ds \\ &\leq h_V \int_0^1 \int_{\gamma_V(0)} \frac{1}{|\gamma_V(s)|} \int_{\gamma_V(s)} u^2(x) dx (1 - s) dy ds \\ &= h_V \int_0^1 \int_{\gamma_V(0)} \frac{1}{|\gamma_V(0)|} \int_{\gamma_V(0)} u^2(F_V(x, s)) dx (1 - s) dy ds \\ &= h_V \int_0^1 \int_{\gamma_V(0)} u^2(F_V(x, s)) dx (1 - s) ds \\ &\simeq \int_{\omega_V} u^2(x) dx. \end{aligned}$$

(ii) To verify the estimate for the  $H^1$ -semi-norm, we rewrite the point-wise gradient:

$$\begin{aligned} (\nabla \Pi^V u)(x) &= \nabla \left( \frac{1}{|\gamma_V(x)|} \int_{\gamma_V(x)} u(y) dy \right) \\ &= \nabla \left( \frac{1}{|\gamma_V(0)|} \int_{\gamma_V(0)} u(F_V(y, \varphi_V(x))) dy \right) \\ &= \frac{1}{|\gamma_V(0)|} \int_{|\gamma_V(0)|} \frac{d(u \circ F_V)}{ds}(y, \varphi_V(x)) \nabla \varphi_V(x) dy \\ &= \frac{1}{|\gamma_V(0)|} \int_{|\gamma_V(0)|} (\nabla u)(F_V(y, \varphi_V(x))) \cdot (V - y) \nabla \varphi_V(x) dy. \end{aligned}$$

Forming the absolute values allows us to estimate

$$\begin{aligned}
|\nabla \Pi^V u|(x) &\leq \frac{1}{|\gamma_V(0)|} \int_{\gamma_V(0)} |(\nabla u)(F_V(y, \varphi_V(x)))| |V - y| |\nabla \varphi_V| dx \\
&\preceq \frac{1}{|\gamma_V(0)|} \int_{\gamma_V(0)} |(\nabla u)(F_V(y, \varphi_V(x)))| h_V h_V^{-1} dy \\
&= \frac{1}{|\gamma_V(x)|} \int_{\gamma_V(x)} |(\nabla u)(y)| dy \\
&= (\Pi^V |\nabla u|)(x).
\end{aligned}$$

The rest follows from the  $L_2$ -estimate (i) applied to  $|\nabla u|$ .

(iii) On the manifold  $\gamma_V(0)$ , there holds the Poincaré inequality

$$\int_{\gamma_V(0)} \left| u(x) - \frac{1}{|\gamma_V(0)|} \int_{\gamma_V(0)} u(y) dy \right|^2 dx \preceq h_V^2 \int_{\gamma_V(0)} |\nabla u|^2 dx.$$

Using  $r_V(x) \simeq (1 - \varphi_V(x))h_V$  we derive

$$\begin{aligned}
&\int_{\omega_V} \frac{1}{r_V^2} (u - \Pi^V u)^2 dx \\
&\simeq h_V \int_0^1 \int_{\gamma_V(0)} \frac{1}{r_V^2} \left( u(F_V(y, s)) - \frac{1}{|\gamma_V(0)|} \int_{\gamma_V(0)} u(F_V(x, s)) dx \right)^2 (1 - s) dy ds \\
&\preceq h_V \int_0^1 \int_{\gamma_V(0)} \frac{h_V^2}{r_V^2} |\nabla_y u(F_V(y, s))|^2 (1 - s) dy ds \\
&\simeq h_V \int_0^1 \int_{\gamma_V(0)} \frac{1}{(1 - s)^2} \left| (\nabla u)(F_V(y, s)) \frac{\partial F_V}{\partial y} \right|^2 (1 - s) dy ds \\
&= h_V \int_0^1 \int_{\gamma_V(0)} |(\nabla u)(F_V(y, s))|^2 (1 - s) dy ds \\
&\simeq \|\nabla u\|_{L_2(\omega_V)}^2.
\end{aligned}$$

(iv) Since  $\varphi_V 1 = \Pi_0^V 1$ , we can subtract the mean value  $\bar{u} := \frac{1}{|\omega_V|} \int_{\omega_V} u(x) dx$ :

$$\begin{aligned}
\|\nabla \{\varphi_V u - \Pi_0^V u\}\| &= \|\nabla \{\varphi_V(u - \bar{u}) - \Pi_0^V(u - \bar{u})\}\| \\
&\leq \|\nabla \{\varphi_V(u - \bar{u})\}\| + \|\nabla \Pi^V(u - \bar{u}) - \Pi^V(u - \bar{u})|_{\gamma_V(0)} \nabla(1 - \varphi_V)\| \\
&\preceq \|(\nabla \varphi_V)(u - \bar{u})\| + \|\varphi_V \nabla u\| + \|\nabla u\| + \left| \Pi^V(u - \bar{u})|_{\gamma_V(0)} \right| \|\nabla \varphi_V\| \\
&\preceq h^{-1} \|u - \bar{u}\| + \|\nabla u\| \\
&\preceq \|\nabla u\|.
\end{aligned}$$

We have used (ii) and the trace inequality for

$$(3) \quad \begin{aligned} \left| \Pi^V(u - \bar{u})|_{\gamma_V(0)} \right| &= \frac{1}{|\gamma_V(0)|} \left| \int_{\gamma_V(0)} u - \bar{u} \, dx \right| \leq \\ &\leq |\gamma_V(0)|^{-1/2} \|u - \bar{u}\|_{L_2(\gamma_V(0))} \leq \|\nabla(u - \bar{u})\| + h^{-1} \|u - \bar{u}\|. \end{aligned}$$

(v) We finally prove  $\|r_V^{-1}\{\varphi_V u - \Pi_0^V u\}\|_{L_2(\omega_V)} \leq \|\nabla u\|_{L_2(\omega_V)}$ . From the definition of  $r_V$ , we get

$$\left\| \frac{1}{r_V} \{\varphi_V u - \Pi_0^V u\} \right\| \simeq \left\| \frac{1}{r_V} \{\varphi_V u - \Pi_0^V u\} \right\| + \sum_{V' \in \omega_V \setminus \{V\}} \left\| \frac{1}{r_{V'}} \{\varphi_{V'} u - \Pi_0^{V'} u\} \right\|.$$

We bound the first term as follows:

$$\begin{aligned} &\left\| \frac{1}{r_V} \{\varphi_V u - \Pi_0^V u\} \right\|_{L_2(\omega_V)} \\ &= \left\| \frac{1}{r_V} \{(1 - (1 - \varphi_V))u - \Pi^V u + (1 - \varphi_V)(\Pi^V u)|_{\gamma_V(0)}\} \right\| \\ &= \left\| \frac{1}{r_V} \{(u - \Pi^V u) - (1 - \varphi_V)(u - \bar{u}) + (1 - \varphi_V)((\Pi^V u)|_{\gamma_V(0)} - \bar{u})\} \right\| \\ &\leq \left\| \frac{1}{r_V} (u - \Pi^V u) \right\| + \left\| \frac{1 - \varphi_V}{r_V} (u - \bar{u}) \right\| + \left\| \frac{1 - \varphi_V}{r_V} ((\Pi^V u)|_{\gamma_V(0)} - \bar{u}) \right\| \\ &\leq \|\nabla u\| + h^{-1} \|u - \bar{u}\| + h^{-1} |(\Pi^V u)|_{\gamma_V(0)} - \bar{u}| |\omega_V|^{1/2} \\ &\leq \|\nabla u\|_{L_2(\omega_V)} \end{aligned}$$

We have used that  $(1 - \varphi_V)/r_V \simeq h^{-1}$ , and applied the Poincaré inequality on  $\omega_V$ , and once again (3).

Before treating the second term, we prove the following estimate on a triangle  $T$ :

$$(4) \quad \int_T \frac{1}{(r_{V'})^2} v^2 \, dx \leq \|\nabla v\|_{L_2(T)}^2,$$

for functions  $v$  vanishing on an edge  $E$  containing the vertex  $V'$ . We transform to a reference triangle  $\hat{T} = \{(x_1, x_2) : 0 \leq x_2 \leq x_1 \leq 1\}$  and use Friedrichs' inequality:

$$\begin{aligned} \int_T \frac{1}{(r_{V'})^2} v^2 \, dx &\simeq h^2 \int_0^1 \int_0^{x_1} \frac{1}{h^2 x_1^2} v^2(x_1, x_2) \, dx_2 \, dx_1 \\ &\leq \int_0^1 \int_0^{x_1} \left( \frac{\partial v}{\partial x_2} \right)^2 \, dx_2 \, dx_1 \leq \|\nabla v\|_{L_2(T)}^2 \end{aligned}$$

Since the function  $v := \varphi_V u - \Pi_0^V u$  vanishes on the boundary  $\partial\omega_V$ , inequality (4) can be applied on each triangle  $T \subset \omega_V$ :

$$\left\| \frac{1}{r_{V'}} \{ \varphi_V u - \Pi_0^V u \} \right\|_{L_2(\omega_V)} \preceq \left\| \nabla \{ \varphi_V u - \Pi_0^V u \} \right\|_{L_2(\omega_V)}$$

Using (iv) and summing over  $V'$ , we get the desired estimate. Due to shape regularity this sum is finite.

This finishes the proof of Lemma 4 □

The *global spider vertex operator* is

$$\Pi_{\mathcal{V}} := \sum_{V \in \mathcal{V}_f} \Pi_0^V.$$

Obviously,  $u - \Pi_{\mathcal{V}} u$  vanishes in any vertex  $V \in \mathcal{V}_f$ . These well-defined zero vertex values are reflected by the following norm definition:

$$(5) \quad \|\cdot\|^2 := \|\nabla \cdot\|_{L_2(\Omega)}^2 + \left\| \frac{1}{r_{\mathcal{V}}} \cdot \right\|_{L_2(\Omega)}^2$$

**Theorem 5.** *Let  $u_1$  be as in Lemma 2. Then, the decomposition*

$$(6) \quad u_1 = \sum_{V \in \mathcal{V}_f} \Pi_0^V u_1 + u_2$$

*is stable in the sense of*

$$(7) \quad \sum_{V \in \mathcal{V}_f} \|\Pi_0^V u_1\|_A^2 + \|u_2\|^2 \preceq \|u\|_A^2.$$

*Proof.* The vertex terms in equation (7) are bounded by

$$\begin{aligned} \|\Pi_0^V u_1\|_A^2 &= \|\Pi^V u_1 - (\Pi^V u_1)|_{\gamma_V(0)}(1 - \varphi_V)\|_A^2 \\ &\leq \|\nabla \Pi^V u_1\|_{L_2(\omega_V)}^2 + |(\Pi^V u_1)|_{\gamma_V(0)}|^2 \|1 - \varphi_V\|_A^2 \\ &\preceq \|\nabla u_1\|_{L_2(\omega_V)}^2 + h_V^{-2} \|u_1\|_{L_2(\omega_V)}^2. \end{aligned}$$

We have used that  $\|1 - \varphi_V\|_A \simeq 1$ . Summing up all terms, one obtains

$$\sum_{V \in \mathcal{V}_f} \|\Pi_0^V u_1\|_A^2 \preceq \|\nabla u_1\|_{L_2(\Omega)}^2 + \|h^{-1} u_1\|_{L_2(\Omega)}^2 \preceq \|u\|_A^2.$$

To bound the second term, we compare with the partition of unity provided by the hat functions:

$$\begin{aligned} \left\| u - \sum_{V \in \mathcal{V}_f} \Pi_0^V u \right\|^2 &= \left\| \sum_{V \in \mathcal{V}_D} \varphi_V u + \sum_{V \in \mathcal{V}_f} (\varphi_V u - \Pi_0^V u) \right\|^2 \\ &\preceq \sum_{V \in \mathcal{V}_D} \|\varphi_V u\|^2 + \sum_{V \in \mathcal{V}_f} \|(\varphi_V u - \Pi_0^V u)\|^2 \end{aligned}$$

The functions  $\varphi_V u$  have a zero-edge for each vertex, and thus, an argument similar to that of Lemma 4, part (v) applies and leads to

$$\|\varphi_V u\|^2 \preceq \|\nabla(\varphi_V u)\|_{L_2(\omega_V)}^2 \leq \|\nabla u\|_{L_2(\omega_V)}^2.$$

□

For the rest of this section,  $u_2$  denotes the second term in the decomposition (6).

**3.3. Edge contributions.** As seen in the last subsection, the remaining function  $u_2$  vanishes in all vertices. We now introduce an edge-based interpolation operator to carry the decomposition further, such that the remaining function,  $u_3$ , contributes only to the inner basis functions of each element.

Therefore we need a lifting operator which extends edge functions to the whole triangle preserving the polynomial order. Such operators were introduced in Babuška et al. [BCM91], and later simplified and extended for 3D by Muñoz-Sola [Mun97]. The lifting on the reference element  $T^R$  with vertices  $(-1, 0)$ ,  $(1, 0)$ ,  $(0, 1)$  and edges  $E_1^R := (-1, 1) \times \{0\}$ ,  $E_2^R$ ,  $E_3^R$  reads:

$$(\mathcal{R}_1 w)(x_1, x_2) := \frac{1}{2x_2} \int_{x_1-x_2}^{x_1+x_2} w(s) ds,$$

for  $w \in L_1([-1, 1])$ . The modification by Muñoz-Sola preserving zero boundary values on the edges  $E_2^R$  and  $E_3^R$  is

$$(\mathcal{R}w)(x_1, x_2) := (1 - x_1 - x_2)(1 + x_1 - x_2) \left( \mathcal{R}_1 \frac{w}{1 - x_1^2} \right)(x_1, x_2).$$

For an arbitrary triangle  $T = F_T(T^R)$  containing the edge  $E = F_T(E_1^R)$ , its transformed version reads

$$\mathcal{R}_T w := \mathcal{R}[w \circ F_T] \circ F_T^{-1}.$$

The Sobolev space  $H_{00}^{1/2}(E)$  on an edge  $E = [V_{E,1}, V_{E,2}]$  is defined by its corresponding norm

$$\|w\|_{H_{00}^{1/2}(E)}^2 := \|w\|_{H^{1/2}(E)}^2 + \int_E \frac{1}{r_{V_E}} w^2 ds,$$

with

$$r_{V_E} := \min\{r_{V_{E,1}}, r_{V_{E,2}}\}.$$

**Lemma 6.** *The Muñoz-Sola lifting operator  $\mathcal{R}_T$  satisfies:*

- (i) *it maps polynomials  $w \in P_0^p(E) := \{v \in P^p(E) : v = 0 \text{ in } V_{E,1}, V_{E,2}\}$  into  $\{v \in P^p(T) : v = 0 \text{ on } \partial T \setminus E\}$ .*
- (ii) *it is bounded in the sense*

$$\|\mathcal{R}_T w\|_{H^1(T)} \preceq \|w\|_{H_{00}^{1/2}(E)}.$$

The proof follows from [BCM91] and [Mun97].

We call  $\omega_E := \omega_{V_{E,1}} \cap \omega_{V_{E,2}}$  the edge patch. We define an *edge-based interpolation operator* as follows:

$$(8) \quad \begin{aligned} \Pi_0^E : \{v \in V_p : v = 0 \text{ in } \mathcal{V}\} &\rightarrow H_0^1(\omega_E) \cap V_p, \\ (\Pi_0^E u)|_T &:= \mathcal{R}_T \operatorname{tr}_E u. \end{aligned}$$

**Lemma 7.** *The edge-based interpolation operator  $\Pi_0^E$  defined in (8) is bounded in the  $\|\cdot\|$ -norm:*

$$\|\nabla \Pi_0^E u\|_{L_2(\omega_E)} \preceq \|u\|_{\omega_E}$$

*Proof.* Foremost, we apply Lemma 6 on a single triangle  $T \subset \omega_E$ :

$$\begin{aligned} \|\nabla \Pi_0^E u\|_{L_2(T)}^2 &= \|\nabla \mathcal{R}_T \operatorname{tr}_E u\|_{L_2(T)}^2 \\ &\preceq \|\operatorname{tr}_E u\|_{H^{1/2}(E)}^2 + \int_E \frac{1}{r_{V_E}} (\operatorname{tr}_E u)^2 ds. \end{aligned}$$

For the first term, the trace theorem can be immediately applied.

The second term, the weighted  $L_2$ -norm on the edge, can be bounded by a weighted norm on the triangle. We transform onto the reference triangle,

$$\int_E \frac{1}{r_{V_E}} u^2 ds = \int_{E_1^R} \frac{1}{r_{V_{E_1^R}}} (u \circ F_T)^2 ds,$$

and write  $u^R := u \circ F_T$ . Due to symmetry, we consider only the right half of the edge  $E_1^R$ , where  $r_{E_1^R} = \frac{1}{1-x_1}$ , and finally apply a trace inequality:

$$\begin{aligned} \int_0^1 \frac{1}{1-x_1} u^R(x_1, 0)^2 dx_1 &\preceq \\ &\preceq \int_0^1 \frac{1}{1-x_1} \int_0^{1-x_1} (1-x_1) \left[ \frac{\partial u^R}{\partial x_2} \right]^2 + \frac{1}{1-x_1} [u^R]^2 dx_2 dx_1 \\ &\preceq \|u^R\|_{T^R}^2 \simeq \|u\|_T^2. \end{aligned}$$

□

This leads us immediately to

**Theorem 8.** *Let  $u_2$  be as in Theorem 5. Then, the decomposition*

$$(9) \quad u_2 = \sum_{E \in \mathcal{E}_f} \Pi_0^E u_2 + u_3$$

*satisfies  $u_3 = 0$  on  $\bigcup_{E \in \mathcal{E}_f} E$  and is bounded in the sense of*

$$(10) \quad \sum_{E \in \mathcal{E}_f} \|\nabla \Pi_0^E u_2\|_{L_2}^2 + \|\nabla u_3\|_{L_2}^2 \preceq \|u_2\|^2.$$

### 3.4. Main result.

*Proof of Theorem 1 for the case of triangles.* Summarizing the last subsections, we have

$$u_1 = u - \Pi_h u, \quad u_2 = u_1 - \sum_{V \in \mathcal{V}_f} \Pi_0^V u_1, \quad u_3 = u_2 - \sum_{E \in \mathcal{E}_f} \Pi_0^E u_2,$$

and the decomposition

$$(11) \quad u = \Pi_h u + \sum_{V \in \mathcal{V}_f} \Pi_0^V u_1 + \sum_{E \in \mathcal{E}_f} \Pi_0^E u_2 + \sum_{T \in \mathcal{T}} u_3|_T.$$

is stable in the  $\|\cdot\|_A$ -norm.

For any edge  $E$  or triangle  $T$ , we can find a vertex  $V$ , such that the corresponding summand is in  $V_V$ . Since for each vertex only finitely many terms appear, we can use the triangle inequality and finally arrive at the missing spectral bound

$$\langle Cu, u \rangle = \inf_{\substack{u = u_0 + \sum_V u_V \\ u_0 \in V_0, u_V \in V_V}} \|u_0\|_A^2 + \sum_V \|u_V\|_A^2 \preceq \langle Au, u \rangle.$$

□

## 4. SUB-SPACE SPLITTING FOR TETRAHEDRA

Most of the proof for the 3D case follows the strategy introduced in Section 3, so we use the definitions thereof. The only principal difference is the edge interpolation operator, which shall be treated in more detail.

**4.1. Coarse and vertex contributions.** We define the level surfaces of the vertex hat basis functions

$$\Gamma_V(x) := \Gamma_V(\varphi_V(x)) := \{y : \varphi_V(y) = \varphi_V(x)\}.$$

As in 2D, we first subtract the coarse grid function

$$u_1 = u - \Pi_h u,$$

and secondly the multi-dimensional vertex interpolant to obtain

$$u_2 = u_1 - \Pi_{\mathcal{V}} u_1,$$

where the definitions of  $\Pi^V$ ,  $\Pi_0^V$ ,  $\Pi_{\mathcal{V}}$  are the same as in Section 3, only the level set lines  $\gamma_V$  are replaced by the level surfaces  $\Gamma_V$ . With the same arguments, one easily shows that

$$(12) \quad \sum_{v \in \mathcal{V}_f} \|\Pi_0^V u_1\|_A^2 + \|\nabla u_2\|_{L_2}^2 + \|r_{\mathcal{V}}^{-1} u_2\|_{L_2}^2 \preceq \|u\|_A^2.$$




FIGURE 3. Reference triangle




FIGURE 4. Edge patch

**4.2. Edge contributions.** Let  $F := \{(s, t) : s \geq 0, t \geq 0, s + t \leq 1\}$  be the reference triangle in Figure 3. For  $(s, t) \in F$ , we define the level lines

$$\gamma_E(s, t) := \{x : \varphi_{V_{E,1}}(x) = s \text{ and } \varphi_{V_{E,2}}(x) = t\},$$

and write

$$\gamma_E(x) := \gamma_E(\varphi_{V_{E,1}}(x), \varphi_{V_{E,2}}(x))$$

for the level line corresponding to a point  $x$  in the edge-patch  $\omega_E$ , see Figure 4.

Define the space of constant functions on these level lines,

$$S_E := \{v : v|_{\gamma_E(x)} = \text{const}\}$$

and its polynomial subspace  $S_{E,p} := S_E \cap V_p$ . The edge averaging operator into  $S_E$  reads

$$(\Pi^E v)(x) := \frac{1}{|\gamma_E(x)|} \int_{\gamma_E(x)} v(y) dy.$$

Furthermore, let  $r_{V_E} := \min\{r_{V_{E,1}}, r_{V_{E,2}}\}$ , and  $r_E(x) := \text{dist}\{x, E\}$ .

**Lemma 9.** *The edge-averaging operator satisfies*

(i)

$$\Pi^E V_p = S_{E,p},$$

(ii)

$$(\Pi^E u)(x) = u(x), \quad \text{for } x \in E, \text{ } u \text{ continuous},$$

(iii)

$$\|\nabla \Pi^E u\|_{L_2(\omega_E)} \leq \|\nabla u\|_{L_2(\omega_E)},$$

(iv)

$$\|r_{V_E}^{-1} \Pi^E u\|_{L_2(\omega_E)} \leq \|r_{V_E}^{-1} u\|_{L_2(\omega_E)},$$

(v)

$$\|r_E^{-1}(u - \Pi^E u)\|_{L_2(\omega_E)} \leq \|\nabla u\|_{L_2(\omega_E)},$$

where  $u \in H^1(\omega_E)$ .

The proof is analogous to the proofs of Lemma 3 and Lemma 4.

Next, the edge-interpolation operator is modified to satisfy zero boundary conditions on  $\partial\omega_E$ . By the isomorphism

$$(13) \quad v_F(s, t) := v|_{\gamma_E(s, t)}, \quad \text{for } v \in S_E,$$

the function space  $S_E$  can be identified with a space on the triangle  $F$ .

**Lemma 10.** *The isomorphism (13) fulfills the following equivalences for functions  $v \in S_E$ :*

$$\begin{aligned} \text{(i)} \quad & \|v\|_{L_2(\omega_E)} \simeq h^{3/2} \|r_{E^R}^{1/2} v_F\|_{L_2(F)}, \\ \text{(ii)} \quad & \|\nabla v\|_{L_2(\omega_E)} \simeq h^{1/2} \|r_{E^R}^{1/2} \nabla v_F\|_{L_2(F)}, \\ \text{(iii)} \quad & \|r_{V_E}^{-1} v\|_{L_2(\omega_E)} \simeq h^{1/2} \left\| \frac{r_{E^R}^{1/2}}{r_{V_{E^R}}} v_F \right\|_{L_2(F)}, \\ \text{(iv)} \quad & \|r_E^{-1} v\|_{L_2(\omega_E)} \simeq h^{1/2} \|r_{E^R}^{-1/2} v_F\|_{L_2(F)}, \end{aligned}$$

where

$$E^R := \{(s, t) \in F : s + t = 1\},$$

$$r_{E^R}(s, t) := 1 - s - t, \quad \text{and} \quad (r_{V_{E^R}})^{-1} := \frac{1}{1-s} + \frac{1}{1-t}.$$

*Proof.* We parameterize the edge-patch  $\omega_E$  by

$$\begin{aligned} F_E : \gamma_E(0, 0) \times F &\rightarrow \omega_E \\ (z, (s, t)) &\mapsto z + s(V_{E,1} - z) + t(V_{E,2} - z). \end{aligned}$$


Note that functions  $v \in S_E$  do not depend on the parameter  $z \in \gamma_E(0, 0)$  and  $v_F(s, t) = (v \circ F_E)(z, s, t)$  for any  $z \in \gamma_E(0, 0)$ . Equivalence (i) holds due to the transformation of the integrals

$$\begin{aligned} \int_{\omega_E} |v|^2 dx &\simeq h^2 \int_{\gamma_E(0,0)} \int_0^1 \int_0^{1-s} |v \circ F_E|^2 (1-s-t) dt ds dz \\ &\simeq h^3 \int_F |v_F|^2 r_{E^R} d(s, t). \end{aligned}$$

Derivatives evaluate to  $\frac{\partial v_F}{\partial s} = (\nabla v) \cdot \frac{\partial F_E}{\partial s} = (\nabla v) \cdot (V_{E,1} - z)$ , and thus

$$|(\nabla v) \circ F_E| \simeq h^{-1} |\nabla v_F|.$$

In combination with (i), we have proven (ii). Finally, equivalences (iii) and (iv) follow from  $r_{V_E} \circ F_E \simeq h r_{V_{E^R}}$ .  $\square$

FIGURE 5. Averaging lines of the smoothing operator  $S_s$ 

We now modify the function

$$(14) \quad u_F(s, t) := (\Pi^E u_2)|_{\gamma_E(s,t)}$$

to obtain a function  $u_{F,00}$  which satisfies zero boundary conditions on the edges  $s = 0$  and  $t = 0$ , and coincides with  $u_F$  on the edge  $s + t = 1$ . This modification is done in such a way that it is continuous in the weighted  $H^1$ -norm.

First, we define the smoothing operator (cf. Figure 5)

$$(S_s v)(s, t) := \int_0^1 v\left(s + \frac{\tau}{2}(1 - s - t), t\right) d\tau.$$

Secondly, we modify the operator to obtain

$$(S_{s,0} v)(s, t) := (S_s v)(s, t) - \frac{1 - s - t}{1 - t} (S_s v)(0, t),$$

which vanishes on the edge  $s = 0$ .

**Lemma 11.** *The smoothing operator  $S_{s,0}$  satisfies*

$$S_{s,0} : \{v \in P^p : v(0, 1) = v(1, 0) = 0\} \rightarrow \{v \in P^p : v(1, 0) = v(0, \cdot) = 0\}$$

and the estimates

$$(15) \quad \begin{aligned} & \left\| r_{ER}^{1/2} \nabla (S_{s,0} v) \right\|_{L_2(F)} + \left\| \frac{r_{ER}^{1/2}}{r_{V_{ER}}} (S_{s,0} v) \right\|_{L_2(F)} + \left\| r_{ER}^{-1/2} (S_{s,0} v - v) \right\|_{L_2(F)} \\ & \preceq \left\| r_{ER}^{1/2} \nabla v \right\|_{L_2(F)} + \left\| \frac{r_{ER}^{1/2}}{r_{V_{ER}}} v \right\|_{L_2(F)}. \end{aligned}$$

*Proof.* Assume that  $v$  is a polynomial vanishing in  $(0, 1)$  and  $(1, 0)$ . Then  $S_s v$  is again a polynomial, which is identical to  $v$  on the edge  $s + t = 1$ . In particular, the restriction onto the edge  $s = 0$  is a polynomial in  $t$  vanishing for  $t = 1$ . Thus, the factor  $1 - t$  in the definition of  $S_{s,0}$  cancels out.

First, we prove the corresponding estimates for the smoothing operator  $S_s$ . We observe that derivatives of  $S_s v$  depend on derivatives of  $v$ , only:

$$\begin{aligned}\frac{\partial(S_s v)}{\partial s} &= \int_0^1 (\nabla v) \cdot \left(1 - \frac{\tau}{2}, 0\right) d\tau, \\ \frac{\partial(S_s v)}{\partial t} &= \int_0^1 (\nabla v) \cdot \left(-\frac{\tau}{2}, 1\right) d\tau.\end{aligned}$$

Since  $r_{ER}$  is bounded from below and from above on the averaging line  $[(s, t); (s + \frac{1}{2}(1 - s - t), t)]$ , the smoothing operator  $S_s$  is bounded in the weighted  $H^1$ -semi-norm. The approximation property corresponding to the weighted  $L_2$ -norm follows from Friedrichs' inequality applied on the same line.

Now, we prove the estimates for the correction  $S_{s,0} - S_s$ . The first is

$$\begin{aligned}& \left\| r_{ER}^{1/2} \nabla \left[ \frac{1-s-t}{1-t} (S_s v)(0, t) \right] \right\|_{L_2(F)} \\ & \leq \left\| r_{ER}^{1/2} \left( \frac{-1}{1-t}, \frac{-s}{(1-t)^2} \right) (S_s v)(0, t) \right\|_{L_2(F)} + \left\| r_{ER}^{1/2} \frac{1-s-t}{1-t} \nabla (S_s v)(0, t) \right\|_{L_2(F)} \\ & \leq \left\| (1-t)^{-1/2} (S_s v)(0, t) \right\|_{L_2(F)} + \left\| (1-t)^{1/2} \frac{\partial(S_s v)}{\partial t}(0, t) \right\|_{L_2(F)} \\ & = \left\| (S_s v)(0, t) \right\|_{L_2(0,1)} + \left\| (1-t) \frac{\partial(S_s v)}{\partial t}(0, t) \right\|_{L_2(0,1)},\end{aligned}$$

the other two are bounded by the same expression.

We now bound these trace norms of  $S_s v$  by the right-hand side of (15). We start with the  $L_2$ -norm:

$$\begin{aligned}\int_0^1 (S_s v)^2(0, t) dt &= \int_0^1 \left[ \int_0^1 v \left( \frac{1-t}{2} \tau, t \right) d\tau \right]^2 dt \\ &\leq \int_0^1 \int_0^1 v^2 \left( \frac{1-t}{2} \tau, t \right) d\tau dt = \int_0^1 \int_0^{\frac{1-t}{2}} v^2(s, t) \frac{2}{1-t} ds dt \\ &\leq \int_F \frac{r_{ER}}{r_{VE}^2} v^2(s, t) ds dt\end{aligned}$$

We have substituted  $s = \frac{1-t}{2} \tau$ , and used that

$$s \leq \frac{1-t}{2} \quad \text{implies} \quad \frac{1}{1-t} \leq 2 \frac{1-s-t}{(1-t)^2} \leq 2 \frac{r_{ER}}{r_{VE}^2}.$$

Similarly, we can bound the weighted  $H^1$ -norm on the edge by

$$\begin{aligned} \int_0^1 (1-t)^2 \left[ \frac{\partial(S_s v)}{\partial t}(0, t) \right]^2 dt &= \int_0^1 (1-t)^2 \left[ \int_0^1 (\nabla v) \cdot (-\tau/2, 1)^T d\tau \right]^2 dt \\ &\preceq \int_0^1 (1-t)^2 \int_0^1 \left| \nabla v \left( \frac{1-t}{2} \tau, t \right) \right|^2 d\tau dt \\ &\preceq \int_0^1 \int_0^{\frac{1-t}{2}} (1-s-t) |\nabla v|^2 ds dt. \end{aligned}$$

□

In the same manner, we define

$$(S_t u)(s, t) := \int_0^1 u \left( s, t + \frac{\tau}{2}(1-s-t) \right) d\tau,$$

and

$$(S_{t,0})(x, y) := (S_t u_{F,0})(s, t) - \frac{1-s-t}{1-s} (S_t u_{F,0})(s, 0).$$

These two smoothing operators allow us to define the function

$$u_{F,00} := S_{t,0} S_{s,0} u_F$$

satisfying zero boundary values at both edges  $s = 0$  and  $t = 0$ .

We define the edge interpolation operator by

$$(16) \quad (\Pi_0^E u_2)(x) := u_{F,00}(\varphi_{V_{E,1}}(x), \varphi_{V_{E,2}}(x)).$$

**Lemma 12.** *The edge interpolation operator  $\Pi_0^E$  satisfies*

$$\|r_{\mathcal{E}}^{-1} \{v - \Pi_0^E v\}\|_{L_2(\omega_E)} \preceq \|\nabla v\|_{L_2(\omega_E)} + \|r_{\mathcal{V}}^{-1} v\|_{L_2(\omega_E)},$$

where

$$r_{\mathcal{E}}(x) := \min_{E \in \mathcal{E}} r_E(x).$$

*Proof.* The proof is analogous to the one of Lemma 4, part (v). We observe that

$$\|r_{\mathcal{E}}^{-1} w\|_{L_2(\omega_E)} \simeq \|r_E^{-1} w\|_{L_2(\omega_E)} + \sum_{E' \subset \omega_E \setminus \{E\}} \|r_{E'}^{-1} w\|_{L_2(\omega_E)}.$$

The desired estimate for the first term follows directly from Lemma 9, Lemma 10 and Lemma 11.

For the second term, we use that  $\int_T r_{E'}^{-2} v^2 dx \preceq \|\nabla v\|_{L_2(T)}^2$  for functions  $v$  vanishing on  $\partial\omega_E$ . □

Finally, we define the *global edge interpolation operator*

$$(17) \quad \Pi_{\mathcal{E}} := \sum_{E \in \mathcal{E}_f} \Pi_0^E,$$

where  $\mathcal{E}_f$  is the set of all free edges, i. e. those which do not lie completely on the Dirichlet boundary. We obtain

**Theorem 13.** *The decomposition*

$$(18) \quad u_2 = \sum_{E \in \mathcal{E}_f} \Pi_0^E u_2 + u_3$$

*fulfills the stability estimate*

$$(19) \quad \sum_{E \in \mathcal{E}_f} \|\Pi_0^E u_2\|_A^2 + \|\nabla u_3\|^2 + \|r_{\mathcal{E}}^{-1} u_3\|^2 \preceq \|\nabla u_2\|^2 + \|r_{\mathcal{V}}^{-1} u_2\|^2.$$

Moreover,  $u_3 = 0$  on  $\bigcup_{E \in \mathcal{E}_f} E$ .

*Proof.* The result is an immediate consequence of Lemma 9, Lemma 11 and Lemma 12 using the argument of finite summation.  $\square$

### 4.3. Main result.

*Proof of Theorem 1 for the case of tetrahedra.* The interpolation on faces in 3D and its analysis follows the line of the edge interpolation in 2D, see Section 3.3.

Summarizing, we obtain

$$\begin{aligned} u_1 &= u - \Pi_h u, & u_2 &= u_1 - \sum_{V \in \mathcal{V}_f} \Pi_0^V u_1, \\ u_3 &= u_2 - \sum_{E \in \mathcal{E}_f} \Pi_0^E u_2, & u_4 &= u_3 - \sum_{F \in \mathcal{F}_f} \Pi_0^F u_3, \end{aligned}$$

where  $\mathcal{F}_f = \{F \in \mathcal{F} : F \not\subset \Gamma_D\}$ . As a consequence of the last subsections, the decomposition

$$(20) \quad u = \Pi_h u + \sum_{V \in \mathcal{V}_f} \Pi_0^V u_1 + \sum_{E \in \mathcal{E}_f} \Pi_0^E u_2 + \sum_{F \in \mathcal{F}_f} \Pi_0^F u_3 + \sum_{T \in \mathcal{T}} u_4|_T$$

is stable in the  $\|\cdot\|_A$ -norm.  $\square$

## 5. NUMERICAL RESULTS

In this section, we show numerical experiments on model problems to verify the theory elaborated in the last sections and to get the absolute condition numbers hidden in the generic constants. Furthermore, we study two more preconditioners.

We consider the  $H^1(\Omega)$  inner product

$$A(u, v) = (\nabla u, \nabla v)_{L_2} + (u, v)_{L_2}$$

on the unit cube  $\Omega = (0, 1)^3$ , which is subdivided into 69 tetrahedra, see Figure 6. We vary the polynomial order  $p$  from 2 up to 10. The condition numbers of the preconditioned systems are computed by the Lanczos method.

**Example 1:** The preconditioner is defined by the space-decomposition with big overlap of Theorem 1:

$$V = V_0 + \sum_{V \in \mathcal{V}} V_V$$

The condition number is proven to be independent of  $h$  and  $p$ . The computed numbers are drawn in Figure 7, labeled 'overlapping V'. The inner unknowns have been eliminated by static condensation. The memory requirement of this preconditioner is considerable: For  $p = 10$ , the memory needed to store the local Cholesky-factors is about 4.4 times larger than the memory required for the global matrix.

In Section 2 we introduced the space splitting into the coarse space  $V_0$  and the vertex subspaces  $V_V$ . However, our proof of Theorem 1 involves the finer splitting of a function  $u$  into a coarse function, functions in the spider spaces  $S_V$ , edge-, face-based and inner functions. Other additive Schwarz preconditioners with uniform condition numbers are induced by this finer splitting.

**Example 2:** Now, we decompose the space into the coarse space, the  $p$ -dimensional spider-vertex spaces  $S_{V,0} = \text{span}\{\varphi_V, \dots, \varphi_V^p\}$ , and the overlapping sub-spaces  $V_E$  on the edge patches:

$$V = V_0 + \sum_{V \in \mathcal{V}} S_{V,0} + \sum_{E \in \mathcal{E}} V_E$$

The condition number is proven to be uniform in  $h$  and  $p$ . The computed values are drawn in Figure 7, labeled 'overlapping E, spider V'. Storing the local factors is now about 80 percent of the memory for the global matrix.

**Example 3:** The interpolation into the spider-vertex space  $S_{V,0}$  has two continuity properties: It is bounded in the energy norm, and the interpolation rest satisfies an error estimate in a weighted  $L_2$ -norm, see Lemma 4 and equation (12). Now, we reduce the  $p$ -dimensional vertex spaces to the spaces spanned by the low energy vertex functions  $\varphi_V^{l.e.}$  defined as solutions of

$$\min_{v \in S_{V,0}, v(V)=1} \|v\|_A^2.$$

These low energy functions can be approximately expressed by the standard vertex functions via  $\varphi_V^{l.e.} = f(\varphi_V)$ , where the polynomial  $f$  solves a weighted 1D problem and can be given explicitly in terms of Jacobi polynomials, see the upcoming report [BPP05]. The interpolation to the low energy vertex space is uniformly bounded, too. But, the approximation estimate in the weighted  $L_2$ -norm depends on  $p$ . The preconditioner is now generated by

$$V = V_0 + \sum_{V \in \mathcal{V}} \text{span}\{\varphi_V^{l.e.}\} + \sum_{E \in \mathcal{E}} V_E.$$

The computed values are drawn in Figure 7, labeled 'overlapping E, low energy V', and show a moderate growth in  $p$ . Low energy vertex basis functions obtained by orthogonalization on the reference element have also been analyzed in [Bic97, SC01].




FIGURE 6. Unstructured mesh




FIGURE 7. Overlapping blocks




FIGURE 8. Standard vertex

**Example 4:** We also tested the preconditioner without additional vertex spaces, i.e.,

$$V = V_0 + \sum_{E \in \mathcal{E}} V_E.$$

Since vertex values must be interpolated by the lowest order functions, the condition number is no longer bounded uniformly in  $p$ . The rapidly growing condition numbers are drawn in Figure 8.

#### REFERENCES

- [Ain96a] M. Ainsworth. Hierarchical domain decomposition preconditioner for  $h$ - $p$  finite element approximation on locally refined meshes. *SIAM J. Sci. Comput.*, 17:1395–1413 (1996)

- [Ain96b] M. Ainsworth. A preconditioner based on domain decomposition for  $h$ - $p$  finite element approximation on quasi-uniform meshes. *SIAM J. Numer. Anal.*, 33:1358–1376 (1996)
- [BS94] I. Babuška and M. Suri. The  $p$  and  $hp$  versions of the finite element method: basic principles and properties. *SIAM Review*, 36:578–632 (1994)
- [BCM91] I. Babuška, A. W. Craig, J. Mandel, and J. Pitkäranta. Efficient preconditioning for the  $p$  version of the finite element method in  $R^2$ . *SIAM J. Numer. Anal.*, 28:624–661 (1991)
- [BPP05] A. Bećirović, P. Paule, V. Pillwein, A. Riese, C. Schneiders, J. Schöberl *Hypergeometric Summation Methods for High Order Finite Elements. Tchn. Report 2005*, SFB013 *Numerical and Symbolic Scientific Computing*, Johannes Kepler University Linz, (2005)
- [BS04] S. Beuchler and J. Schöberl. Extension operators on tensor product structures in 2D and 3D. *Applied Numerical Mathematics* (available online Nov 2004)
- [BSS04] S. Beuchler, R. Schneider, and Ch. Schwab. Multiresolution weighted norm equivalences and applications. *Numerische Mathematik*, 98:67–97, (2004)
- [Bic97] I. Bica. *Iterative Substructuring Algorithms for the  $p$ -version Finite Element Method for Elliptic Problems*. PhD thesis, Courant Institute of Mathematical Sciences, New York University (1997)
- [Cas97] M. Casarin. Quasi-optimal Schwarz methods for the conforming spectral element discretization. *SIAM J. Numer. Anal.*, 34:2482–2502 (1997)
- [Cle75] P. Clément. Approximation by finite element functions using local regularization. *R.A.I.R.O. Anal. Numer.*, pages 77–84 (1975)
- [DW90] M. Dryja and O. B. Widlund. Towards a unified theory of domain decomposition algorithms for elliptic problems. In T. F. Chan, R. Glowinski, J. Périaux, and O. B. Widlund, editors, *Third International Symposium on Domain Decomposition Methods for Partial Differential Equations*, pages 3–21, Philadelphia, (1990)
- [EM04] T. Eibner and J. M. Melenk A local error analysis of the boundary concentrated FEM. *IMA J. Numer. Anal.*, (to appear)
- [GO95] M. Griebel and P. Oswald. On the abstract theory of additive and multiplicative Schwarz algorithms. *Numer. Math.*, 70:163–180 (1995)
- [GC98] B. Guo and W. Cao. An additive Schwarz method for the  $h$ - $p$  version of the finite element method in three dimensions. *SIAM J. Numer. Anal.*, 35:632–654 (1998)
- [KS99] G. Karniadakis and S. Sherwin “Spectral/hp Element Methods for CFD”, Oxford University Press, Oxford (1999)
- [KJ99] V. Korneev and S. Jensen. Domain decomposition preconditioning in the hierarchical  $p$ -version of the finite element method. *Appl. Numer. Math.*, 29:479–518 (1999)
- [Man90] J. Mandel. Iterative solvers by substructuring for the  $p$ -version finite element method. *Comput. Methods. Appl. Mech. Eng.*, 80:117–128 (1990)
- [Mel02] J. M. Melenk. On condition numbers in  $hp$ -FEM with Gauss-Lobatto based shape functions. *J. Comp. Appl. Math.*, 139:21–48 (2002)
- [Mun97] R. Muñoz-Sola. Polynomial liftings on a tetrahedron and applications to the  $hp$ -version of the finite element method in three dimensions. *SIAM J. Numer. Anal.*, 34(1):282–314 (1997)
- [Pav94] L. Pavarino. Additive Schwarz methods for the  $p$ -version finite element method. *Numerische Mathematik*, 66:493–515 (1994)
- [SMPZ05] J. Schöberl, J. M. Melenk, C. G. A. Pechstein, and S. C. Zanglmayr. Schwarz Preconditioning for High Order Simplicial Finite Elements. proc. to 16<sup>th</sup> *International Conference on Domain Decomposition Methods*, New York, 2005, ed. D. Keyes and O. Widlund (submitted, 12 pages)
- [Sch98] C. Schwab.  *$p$ - and  $hp$ -Finite Element Methods. Theory and Applications in Solid and Fluid Mechanics*. Oxford Science Publications (1998)

- [SC01] S. J. Sherwin and M. Casarin. Low Energy Basis Preconditioning for Elliptic Substructured Solvers Based on Unstructured Spectral/ hp Element Discretisations. *Journal of Computational Physics*, 171:1–24 (2001)
- [SBG96] B. F. Smith, P. E. Bjørstad, and W. D. Gropp. *Domain Decomposition. Parallel Multilevel Methods for Elliptic Partial Differential Equations*. Cambridge University Press (1996)
- [SB91] B. Szabó and I. Babuška. *Finite Element Analysis*. Wiley, (1991)
- [SDR04] B. Szabó, A. Düster, E. Rank. *The p-version of the finite element method*. In E. Stein, R de Borst, T.J.R. Hughes, eds, *Encyclopedia of Computational Mechanics*, John Wiley & Sons, (2004)
- [TW04] A. Toselli and O. Widlund. *Domain Decomposition Methods - Algorithms and Theory*. Springer Series in Computational Mathematics, Vol. 34 (2005)
- [Qua99] A. Quarteroni and A. Valli *Domain Decomposition Methods for Partial Differential Equations*. Oxford University Press, Oxford (1999)

J. SCHÖBERL, JOHANN RADON INSTITUTE FOR COMPUTATIONAL AND APPLIED MATHEMATICS (RICAM), AUSTRIA, [JOACHIM.SCHOEBERL@OEAW.AC.AT](mailto:JOACHIM.SCHOEBERL@OEAW.AC.AT)

J. M. MELENK, THE UNIVERSITY OF READING, DEPARTMENT OF MATHEMATICS, UK, [J.M.MELENK@READING.AC.UK](mailto:J.M.MELENK@READING.AC.UK)

C. G. A. PECHSTEIN, INSTITUTE FOR COMPUTATIONAL MATHEMATICS, JOHANNES KEPLER UNIVERSITY LINZ, AUSTRIA, [CLEMENS.PECHSTEIN@NUMA.UNI-LINZ.AC.AT](mailto:CLEMENS.PECHSTEIN@NUMA.UNI-LINZ.AC.AT)

S. C. ZAGLMAYR, JOHANN RADON INSTITUTE FOR COMPUTATIONAL AND APPLIED MATHEMATICS (RICAM), AUSTRIA, [SABINE.ZAGLMAYR@OEAW.AC.AT](mailto:SABINE.ZAGLMAYR@OEAW.AC.AT)