

Three-Dimensional Symmetric Shift Radix Systems

**A. Huszti, K. Scheicher, P. Surer, J.M.
Thuswaldner**

RICAM-Report 2007-06

THREE-DIMENSIONAL SYMMETRIC SHIFT RADIX SYSTEMS

ANDREA HUSZTI, KLAUS SCHEICHER, PAUL SURER, AND JÖRG M. THUSWALDNER

ABSTRACT. Shift radix systems have been introduced by Akiyama *et al.* as a common generalization of β -expansions and canonical number systems. In the present paper we study a variant of them, so-called symmetric shift radix systems which were introduced recently by Akiyama and Scheicher. In particular, for $d \in \mathbb{N}$ and $\mathbf{r} \in \mathbb{R}^d$ let

$$\tau_{\mathbf{r}} : \mathbb{Z}^d \rightarrow \mathbb{Z}^d, \quad \mathbf{a} = (a_1, \dots, a_d) \mapsto \left(a_2, \dots, a_d, - \left\lfloor r_1 a_1 + r_2 a_2 + \dots + r_d a_d + \frac{1}{2} \right\rfloor \right).$$

The mapping $\tau_{\mathbf{r}}$ is called a *symmetric shift radix system*, if

$$\forall \mathbf{a} \in \mathbb{Z}^d \quad \exists n \in \mathbb{N} : \tau_{\mathbf{r}}^n(\mathbf{a}) = \mathbf{0}.$$

Akiyama and Scheicher found out that the parameters \mathbf{r} giving rise to a symmetric shift radix system in \mathbb{R}^2 form an isosceles triangle together with parts of its boundary. In the present paper we completely characterize all symmetric shift radix systems in the three dimensional space. The result is that $\mathbf{r} \in \mathbb{R}^3$ gives rise to a symmetric shift radix system $\tau_{\mathbf{r}}$ if and only if \mathbf{r} is contained in the union of three convex polyhedra (together with some parts of their boundary). We describe this set explicitly.

1. INTRODUCTION

In Akiyama *et al.* [1] a dynamical system called shift radix system has been introduced.

Definition 1.1 (*cf.* [1]). Let $d \geq 1$ be an integer, $\mathbf{r} \in \mathbb{R}^d$, and let

$$\tilde{\tau}_{\mathbf{r}} : \mathbb{Z}^d \rightarrow \mathbb{Z}^d, \quad \mathbf{a} = (a_1, \dots, a_d) \mapsto (a_2, \dots, a_d, -\lfloor \mathbf{r}\mathbf{a} \rfloor),$$

where $\mathbf{r}\mathbf{a} = r_1 a_1 + r_2 a_2 + \dots + r_d a_d$, *i.e.*, the inner product of the vectors \mathbf{r} and \mathbf{a} . Then $\tilde{\tau}_{\mathbf{r}}$ is called a *shift radix system* (for short SRS), if

$$\forall \mathbf{a} \in \mathbb{Z}^d \quad \exists n \in \mathbb{N} : \tilde{\tau}_{\mathbf{r}}^n(\mathbf{a}) = \mathbf{0}.$$

SRS are related to number systems as β -expansions (*cf.* for instance [8, 10, 13]) or canonical number systems (*cf.* for instance [12]). Indeed they form a unification and generalization of these notions of number systems. More details about SRS and their relation to β -expansions and CNS can be found in [1, 3, 4, 15]. In this paper we want to deal with a variant of SRS, the so-called symmetric shift radix systems.

Definition 1.2 (*cf.* [5]). Let $d \geq 1$ be an integer, $\mathbf{r} \in \mathbb{R}^d$, and let

$$(1.1) \quad \tau_{\mathbf{r}} : \mathbb{Z}^d \rightarrow \mathbb{Z}^d, \quad \mathbf{a} = (a_1, \dots, a_d) \mapsto \left(a_2, \dots, a_d, - \left\lfloor \mathbf{r}\mathbf{a} + \frac{1}{2} \right\rfloor \right).$$

Then $\tau_{\mathbf{r}}$ is called a *symmetric shift radix system* (SSRS for short), if

$$\forall \mathbf{a} \in \mathbb{Z}^d \quad \exists n \in \mathbb{N} : \tau_{\mathbf{r}}^n(\mathbf{a}) = \mathbf{0}.$$

Observe that the only difference between the two definitions is just the additional summand “ $+\frac{1}{2}$ ” inside the floor function in (1.1).

Date: August 22, 2006.

2000 Mathematics Subject Classification. 11A63.

Key words and phrases. beta expansion, canonical number system, shift radix system.

The third and fourth author are supported by the Austrian Research Foundation (FWF), Project S9610, that is part of the Austrian Research Network “Analytic Combinatorics and Probabilistic Number Theory”.

All authors are supported by the “Aktion Österreich-Ungarn” project 63öu3.

SSRS have been already treated by Akiyama and Scheicher [5]. It was proved there that, analogously to the classical SRS, we have a strong relationship to certain notions of number systems. In particular SSRS form a common generalization of symmetric β -expansions and symmetric canonical number systems (SCNS). For the sake of completeness we recall the definition of these number systems and summarize the results on their relation to SSRS.

Definition 1.3 (cf. [5]). Let $\beta > 1$ be a real non-integral number. The unique representation of a positive $\gamma \in \mathbb{R}$ of the form

$$\gamma = d_m \beta^m + d_{m-1} \beta^{m-1} + d_{m-2} \beta^{m-2} + \dots$$

for some $m \in \mathbb{Z}$ with $d_k \in (-\frac{\beta+1}{2}, \dots, \frac{\beta+1}{2}) \cap \mathbb{Z}$, $k \leq m$, such that the condition

$$-\frac{\beta^{k+1}}{2} \leq \sum_{i \leq k} d_i \beta^i < \frac{\beta^{k+1}}{2}$$

is satisfied for any $k \leq m$, is called the *symmetric β -expansion of γ* . We say that β has property (SF) if all $\gamma \in \mathbb{Z}[\beta^{-1}]$ admit a finite symmetric β -expansion.

In the same way as for property (F) of ordinary β -expansions (see [8]) it can be shown that a number β with property (SF) is necessarily a Pisot number.

Theorem 1.4 (cf. [5, Theorem 3.6]). *A Pisot number β with minimal polynomial $(x - \beta)(x^{d-1} + r_{d-1}x^{d-2} + \dots + r_2x + r_1)$ has Property (SF) if and only if $\tau_{(r_1, \dots, r_{d-1})}$ is an SSRS.*

There is a similar statement for SCNS whose definition we want to recall now.

Definition 1.5 (cf. [5]). Let $P(x) = x^d + a_{d-1}x^{d-1} + \dots + a_1x + a_0 \in \mathbb{Z}[x]$, $|a_0| \geq 2$, $\mathcal{R} := \mathbb{Z}[x]/P(x)\mathbb{Z}[x]$, $X \in \mathcal{R}$ the image of x under the canonical epimorphism from $\mathbb{Z}[x]$ to \mathcal{R} and $\mathcal{N} := \left[-\frac{|a_0|}{2}, \frac{|a_0|}{2} \right) \cap \mathbb{Z}$. $(P(x), \mathcal{N})$ is called a *symmetric canonical number system* (SCNS) if each $R \in \mathcal{R}$ can be represented as

$$R = \sum_{i=0}^n l_i X^i, \quad l_i \in \mathcal{N}.$$

Theorem 1.6 (cf. [5, Theorem 2.1]). *$(P(x), \mathcal{N})$ with $P(x) = x^d + a_{d-1}x^{d-1} + \dots + a_1x + a_0 \in \mathbb{Z}[x]$ and $\mathcal{N} := \left[-\frac{|a_0|}{2}, \frac{|a_0|}{2} \right) \cap \mathbb{Z}$ is an SCNS if and only if $\tau_{\mathbf{r}}$ is an SSRS, where $\mathbf{r} = \left(\frac{1}{a_0}, \frac{a_{d-1}}{a_0}, \dots, \frac{a_1}{a_0} \right)$.*

Now, in order to show the differences between SSRS and SRS, we define the following sets related to the behavior of the periods of $\tilde{\tau}_{\mathbf{r}}$ and $\tau_{\mathbf{r}}$, respectively. Let

$$\begin{aligned} \tilde{\mathcal{D}}_d &:= \{ \mathbf{r} \in \mathbb{R}^d \mid \forall \mathbf{a} \in \mathbb{Z}^d \exists n, l \in \mathbb{N} : \tilde{\tau}_{\mathbf{r}}^k(\mathbf{a}) = \tilde{\tau}_{\mathbf{r}}^{k+l}(\mathbf{a}) \forall k \geq n \} \text{ and} \\ \tilde{\mathcal{D}}_d^0 &:= \{ \mathbf{r} \in \mathbb{R}^d \mid \tilde{\tau}_{\mathbf{r}} \text{ is an SRS} \}, \end{aligned}$$

as well as


$$\begin{aligned} \mathcal{D}_d &:= \{ \mathbf{r} \in \mathbb{R}^d \mid \forall \mathbf{a} \in \mathbb{Z}^d \exists n, l \in \mathbb{N} : \tau_{\mathbf{r}}^k(\mathbf{a}) = \tau_{\mathbf{r}}^{k+l}(\mathbf{a}) \forall k \geq n \} \text{ and} \\ \mathcal{D}_d^0 &:= \{ \mathbf{r} \in \mathbb{R}^d \mid \tau_{\mathbf{r}} \text{ is an SSRS} \}. \end{aligned}$$

For $\mathbf{r} = (r_1, \dots, r_d) \in \mathbb{R}^d$, let

$$R(\mathbf{r}) = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ \vdots & 0 & \ddots & \ddots & \vdots \\ \vdots & \vdots & \ddots & 1 & 0 \\ 0 & 0 & \dots & 0 & 1 \\ -r_1 & -r_2 & \dots & -r_{d-1} & -r_d \end{pmatrix}.$$

For $M \in \mathbb{R}^{d \times d}$, denote by $\varrho(M)$ the spectral radius of M , i.e., the maximum absolute value of the eigenvalues of M . For simplicity, we write $\varrho(\mathbf{r}) := \varrho(R(\mathbf{r}))$. Let

$$\mathcal{E}_d(\varepsilon) = \{ \mathbf{r} \in \mathbb{R}^d : \varrho(\mathbf{r}) < \varepsilon \}.$$

FIGURE 1. An approximation of $\tilde{\mathcal{D}}_2^0$

It is known that the $\overline{\mathcal{E}_d(\varepsilon)}$ is a regular set, *i.e.*, the set coincides with the closure of its interior.

We start with the comparison of the sets \mathcal{D}_d and $\tilde{\mathcal{D}}_d$. Firstly, it can easily be seen that their interiors are the same since from [1] we know $\mathcal{E}_d(1) \subset \tilde{\mathcal{D}}_d \subset \overline{\mathcal{E}_d(1)}$ while in [5] it has been shown that

$$(1.2) \quad \mathcal{E}_d(1) \subset \mathcal{D}_d \subset \overline{\mathcal{E}_d(1)}.$$

We will dwell upon the set \mathcal{D}_d in Section 2. However, the sets \mathcal{D}_d^0 and $\tilde{\mathcal{D}}_d^0$ have different behavior. Properties of the set $\tilde{\mathcal{D}}_d^0$ have been developed in [1, 3, 4]. In [3, 15] special attention was paid to the two dimensional case $\tilde{\mathcal{D}}_2^0$. It turns out that the structure of $\tilde{\mathcal{D}}_2^0$ is very complicated and despite large parts of the set could be characterized, a full characterization is still outstanding. An approximation of $\tilde{\mathcal{D}}_2^0$ is shown in Figure 1.

The sets $\tilde{\mathcal{D}}_d^0$ for $d \geq 3$ are not yet investigated in detail, however, computer experiments indicate that $\tilde{\mathcal{D}}_3^0$ is hard to describe.

For the case of SSRS the situation becomes more pleasant at least for low dimensions. Akiyama and Scheicher [5] presented the surprising result that \mathcal{D}_2^0 has a really simple characterization (see Figure 2). They found out that

$$\mathcal{D}_2^0 = \left\{ (x, y) \in \mathbb{R}^2 \mid x \leq \frac{1}{2}, -x - \frac{1}{2} < y \leq x + \frac{1}{2} \right\} \setminus \left\{ \left(\frac{1}{2}, y \right) \in \mathbb{R}^2 \mid \frac{1}{2} < y < 1 \right\},$$


i.e., \mathcal{D}_2^0 is an isosceles triangle together with some parts of its boundary. In the present paper we are interested in the shape of the set \mathcal{D}_3^0 . Amazingly, we will see that \mathcal{D}_3^0 can be described completely as a simple as well as interesting body.

The paper is organized as follows. In Section 2 we concentrate on \mathcal{D}_d and its relation with \mathcal{D}_d^0 in general and specially if $d = 3$. Furthermore, we present an algorithm that is useful for the description of \mathcal{D}_d^0 . It was firstly presented in [6] and in progression has been adapted for SSRS in [5]. In Section 3 we will state the exact characterization of the set \mathcal{D}_3^0 . In Section 4 we will prove this characterization result by using the algorithm presented in Section 2 together with some other algorithms related to bodies defined by polynomial inequalities such as the cylindrical algebraic decomposition algorithm (*cf.* Collins [7]).

2. CONSTRUCTION OF \mathcal{D}_3^0 FROM \mathcal{D}_3

Let us consider the set \mathcal{D}_d . By (1.2) apart from the boundary, the set \mathcal{D}_d coincides with the set $\mathcal{E}_d(1)$ and their closures are equal. As the minimal polynomial of $R(\mathbf{r})$ is given by

$$(2.1) \quad x^d + r_d x^{d-1} + \dots + r_2 x + r_1$$

FIGURE 2. The shape of \mathcal{D}_2^0

the problem of characterizing $\mathcal{E}_d(\varepsilon)$ is equivalent to the problem of finding polynomials of the form (2.1) whose roots lie inside the ε multiple of the unit ball. This problem was already settled in [14, 17]. From these references we easily get the following lemma.

Lemma 2.1. *A vector $\mathbf{r} = (r_1, \dots, r_d)$ is contained in $\mathcal{E}_d(\varepsilon)$ if and only if the Hermitian form*

$$H_d(x_0, \dots, x_{d-1}) := \sum_{i=0}^{d-1} \left| \sum_{j=i}^{d-1} \varepsilon^{d+i-j} r_{d+i-j+1} x_j \right|^2 - \sum_{i=0}^{d-1} \left| \sum_{j=i}^{d-1} \varepsilon^{j-i} r_{j-i+1} x_j \right|^2$$

with $r_{d+1} = 1$ is positive definite.

Now we turn to the study of \mathcal{D}_d^0 . The set \mathcal{D}_d^0 can be constructed from the set \mathcal{D}_d by cutting out convex polyhedra. For $\mathbf{r} = (r_1, \dots, r_d) \in \mathcal{D}_d$, an element $\mathbf{a} = (a_1, \dots, a_d) \in \mathbb{Z}^d \setminus \{0\}$ is a non-zero periodic point of $\tau_{\mathbf{r}}$ of length L , if $\mathbf{a} = \tau_{\mathbf{r}}^L(\mathbf{a})$. From the definition of \mathcal{D}_d^0 it follows that the existence of such a periodic point is necessary and sufficient for $\mathbf{r} \notin \mathcal{D}_d^0$. Suppose that the period defined by \mathbf{a} runs through the points

$$\tau_{\mathbf{r}}^j(\mathbf{a}) = (a_{1+j}, \dots, a_{d+j}) \quad (0 \leq j \leq L-1),$$

where $a_{L+1} = a_1, \dots, a_{L+d-1} = a_{d-1}$. We denote such a period by $(a_1, \dots, a_d); a_{d+1}, \dots, a_L$ and say that it is a period of $\tau_{\mathbf{r}}$ or just a period of \mathcal{D}_d .

Let a non-zero period $\pi := (a_1, \dots, a_d); a_{d+1}, \dots, a_L$ be given. We may ask for the set $P(\pi)$ of all $\mathbf{r} \in \mathcal{D}_d$ for that π occurs as a period of $\tau_{\mathbf{r}}$. By of the definition of $\tau_{\mathbf{r}}$, an element $\mathbf{r} \in P(\pi)$ has to satisfy the system of L double inequalities

$$(2.2) \quad -\frac{1}{2} \leq r_1 a_{1+i} + r_2 a_{2+i} + \dots + r_d a_{d+i} + a_{d+1+i} < \frac{1}{2}.$$

Here i runs from 0 to $L-1$ and $a_{L+1} = a_1, \dots, a_{L+d} = a_d$. Such a system characterizes a convex polyhedron, which is possibly degenerated or equal to the empty set. Therefore we will call $P(\pi)$ a *cutout polyhedron*. Example 2.3 shows how $P(\pi)$ could look like for a given period in the three dimensional case. Since each point $\mathbf{r} \in P(\pi)$ has π as a period of the associated mapping $\tau_{\mathbf{r}}$ the set $P(\pi)$ has empty intersection with \mathcal{D}_d^0 . Thus we get the representation

$$\mathcal{D}_d^0 = \mathcal{D}_d \setminus \bigcup_{\pi \neq 0} P(\pi),$$

where the union is extended over all non-zero periods π . Since the set of periods is infinite, this expression is not suitable for calculations. The following theorem shows how to reduce the set of possible periods to a finite set and gives an efficient algorithm for a closed subset H of $\text{int } \mathcal{D}_d = \mathcal{E}_d(1)$ to determine $H \cap \mathcal{D}_d^0$. It was presented for the first time for CNS in [6] and further improved and adapted to SRS in [1, 3, 15]. In [5] the algorithm was established for SSRS. Basically

we will use this version. Let \mathbf{e}_i be the i -th canonical unit vector. For an $\mathbf{r} = (r_1, \dots, r_d) \in \text{int } \mathcal{D}_d$, denote by $\mathcal{V}(\mathbf{r}) \subset \mathbb{Z}^d$ the smallest set with the properties

- (1) $\pm \mathbf{e}_i \in \mathcal{V}(\mathbf{r}), i = 1, \dots, d,$
- (2) $(a_1, \dots, a_d) \in \mathcal{V}(\mathbf{r}) \Rightarrow (a_2, \dots, a_{d+1}) \in \mathcal{V}(\mathbf{r})$ where a_{d+1} satisfies

$$-1 < r_1 a_1 + r_2 a_2 + \dots + r_d a_d + a_{d+1} < 1.$$

$\mathcal{V}(\mathbf{r}) \subset \mathbb{Z}^d$ is called a *set of witnesses* for \mathbf{r} . Additionally define $\mathcal{G}(\mathcal{V}(\mathbf{r})) = V \times E$ to be the graph with set of vertices $V = \mathcal{V}(\mathbf{r})$ and set of edges $E \subset V \times V$ such that

$$\forall \mathbf{a} \in V : (\mathbf{a}, \tau_{\mathbf{r}}(\mathbf{a})) \in E.$$

The set of vertices is exactly the same as in [1]. The edges are defined in a different way. There exists only one outgoing edge for each vertex. We are interested in the cyclic structure of such graphs. A cycle $\mathbf{a}_1 \rightarrow \mathbf{a}_2 \rightarrow \dots \rightarrow \mathbf{a}_L \rightarrow \mathbf{a}_1$ induces a period of length L in an obvious way.

Theorem 2.2 (cf. [5, Theorem 4.2]). *Let $\mathbf{r}_1, \dots, \mathbf{r}_k \in \mathcal{D}_d$ and let $H := \square(\mathbf{r}_1, \dots, \mathbf{r}_k)$ be the convex hull of $\mathbf{r}_1, \dots, \mathbf{r}_k$. Assume that $H \subset \text{int } \mathcal{D}_d$ and sufficiently small in diameter. Then there exists an algorithm to construct a finite directed graph $G(H) = V \times E$ with vertices $V \subset \mathbb{Z}^d$ and edges $E \subset V \times V$ which satisfies*

- (1) $\pm \mathbf{e}_i \in V$ for all $i = 1, \dots, d,$
- (2) $\mathcal{G}(\mathcal{V}(\mathbf{x}))$ is a subgraph of $G(H)$ for all $\mathbf{x} \in H,$
- (3) $H \cap \mathcal{D}_d^0 = H \setminus \bigcup_{\pi} P(\pi)$, where π are taken over all periods induced by the nonzero primitive cycles of G .

Observe that the theorem can be extended to any convex set $H \subset \text{int } \mathcal{D}_d$ analogously to [15]. In our context the version presented in Theorem 2.2 suffices. In practice, the graph in Theorem 2.2 is constructed by successively adding new vertices. Note that the restriction ‘‘sufficiently small’’ is not superfluous. It turns out that the size of the set of vertices in the graph in Theorem 2.2 can grow to infinity if H is chosen too big. For more detailed information on this topic, see [5, 15]. For us it is only important to choose H in a way that everything stays finite. This can be realized by a suitable subdivision of a given set. We will turn to this problem in Section 4.

Theorem 2.2 proved to be a powerful tool for characterizing \mathcal{D}_d^0 . If it is used properly, $\mathcal{D}_d^0 \cap H$ can be characterized for any closed $H \subset \text{int } \mathcal{D}_d$. Thus, whenever there exists such an H with $\mathcal{D}_d^0 \subset H$ there is a chance to characterize \mathcal{D}_d^0 completely. That was the case for $d = 2$ and we will see that this is valid for $d = 3$, too. For classical SRS, there does not exist such a set H for $d \geq 2$.

Our aim is to characterize \mathcal{D}_3^0 . We already know that

$$\mathcal{E}_3(1) \subset \mathcal{D}_3 \subset \overline{\mathcal{E}_3(1)}.$$

From Lemma 2.1 we calculate

$$\mathcal{E}_3(1) = \{(x, y, z) \in \mathbb{R}^3 \mid |x| < 1, |y - xz| < 1 - x^2, |x + z| < |y + 1|\}.$$

The following example shows how a given period cuts out a polyhedron from $\mathcal{E}_3(1)$.


Example 2.3. Consider the period $\pi := (1, 1, -1); -1, 0$. It induces a system of inequalities (2.2) which describes the polyhedron $P(\pi)$. In our case we get

$$P(\pi) = \left\{ (x, y, z) \mid -\frac{1}{2} \leq x + y - z - 1 < \frac{1}{2} \wedge -\frac{1}{2} \leq x - y - z < \frac{1}{2} \wedge -\frac{1}{2} \leq -x - y + 1 < \frac{1}{2} \right. \\ \left. \wedge -\frac{1}{2} \leq -x + z + 1 < \frac{1}{2} \wedge -\frac{1}{2} \leq y + z - 1 < \frac{1}{2} \right\}.$$

By removing redundant inequalities, this reduces to

$$P(\pi) = \left\{ (x, y, z) \mid x + y - z - 1 < \frac{1}{2} \wedge x - y - z < \frac{1}{2} \wedge -\frac{1}{2} \leq -x - y + 1 \right. \\ \left. \wedge -x + z + 1 < \frac{1}{2} \wedge -\frac{1}{2} \leq y + z - 1 \right\}$$

yielding a polyhedron with five faces. $P(\pi)$ only contains $\mathbf{r} \in \mathcal{D}_d$ with $\tau_{\mathbf{r}}^5((1, 1, -1)) = (1, 1, -1)$ and, hence, $P(\pi)$ has empty intersection with \mathcal{D}_3^0 . Figure 3 shows the position of $P(\pi)$ in $\mathcal{E}_3(1)$. It is easy to see that $P(\pi)$ really cuts out something.

FIGURE 3. The position of $P(\pi)$ in $\mathcal{E}_3(1)$

In the sequel we will need $\overline{\mathcal{E}_3(1)}$ and there some problems occur. Suppose the set which is obtained by changing all the strict inequalities of $\mathcal{E}_3(1)$ to not strict ones. One may think that it equals $\overline{\mathcal{E}_3(1)}$, but this is not the case. It can be easily seen that this set contains the unbounded lines $(1, \lambda, \lambda), \lambda \in \mathbb{R}$ and $(-1, \mu, -\mu), \mu \in \mathbb{R}$ which cannot be true for $\overline{\mathcal{E}_3(1)}$. Hence, $\overline{\mathcal{E}_3(1)}$ is only a subset of this set. We will solve the problem by adding some suitable inequalities. Let

$$\mathcal{E}'_3 := \{(x, y, z) \in \mathbb{R}^3 \mid |x| \leq 1, |y - xz| \leq 1 - x^2, |x + z| \leq |y + 1| \wedge |y - 1| \leq 2 \wedge |z| \leq 3\}$$

and consider the intersection of \mathcal{E}'_3 with the hyperplane $A_c := \{(x, y, z) \in \mathbb{R}^3 \mid x - c = 0\}$ for constant c .

Lemma 2.4. *For any $|c| < 1$ the intersection of \mathcal{E}'_3 with the plane A_c yields the closed triangle $\Delta(A_c^{(1)}, A_c^{(2)}, A_c^{(3)})$ with $A_c^{(1)} = (c, -1, -c), A_c^{(2)} = (c, 1 - 2c, c - 2), A_c^{(3)} = (c, 2c + 1, c + 2)$.*

Proof. We have

$$\mathcal{E}'_3 \cap A_c = \{(c, y, z) \in \mathbb{R}^3 \mid |y - cz| \leq 1 - c^2, |c + z| \leq |y + 1| \wedge |y - 1| \leq 2 \wedge |z| \leq 3\}.$$

As all inequalities are linear, this is a convex set. It is quickly verified that $A_c^{(1)}, A_c^{(2)}, A_c^{(3)} \in \mathcal{E}'_3 \cap A_c$. Thus $\Delta(A_c^{(1)}, A_c^{(2)}, A_c^{(3)}) \subset \mathcal{E}'_3 \cap A_c$. On the other hand consider the closed convex set

$$B_c := \{(c, y, z) \mid |y - cz| \leq 1 - c^2 \wedge c + z \leq y + 1 \wedge -y - 1 \leq c + z\}.$$

Observe that for its definition we used only inequalities that occurred in the definition of $\mathcal{E}'_3 \cap A_c$ and hence we have $\mathcal{E}'_3 \cap A_c \subset B_c$. Pairwise intersection of the three boundary lines of B_c yields exactly the three points $A_c^{(1)}, A_c^{(2)}, A_c^{(3)}$ and therefore $\Delta(A_c^{(1)}, A_c^{(2)}, A_c^{(3)}) = B_c \supset \mathcal{E}'_3 \cap A_c$. \square

Theorem 2.5. $\overline{\mathcal{E}_3(1)} = \mathcal{E}'_3$.

Proof. Obviously \mathcal{E}'_3 is a closed set while $\mathcal{E}_3(1)$ is open. We state that $\text{int } \mathcal{E}'_3 = \mathcal{E}_3(1)$. From Lemma 2.4 we know

$$\mathcal{E}'_3 \cap A_c = \{(c, y, z) \mid |y - cz| \leq 1 - c^2 \wedge c + z \leq y + 1 \wedge -y - 1 \leq c + z\}$$

and as every point of $\mathcal{E}_3(1)$ is inside $\mathcal{E}'_3 \cap A_c$ for some $|c| < 1$ we have

$$\mathcal{E}'_3 = \bigcup_{|c| < 1} (\mathcal{E}'_3 \cap A_c) \supset \mathcal{E}_3(1)$$

and therefore

$$\text{int } \mathcal{E}'_3 \supset \text{int } \mathcal{E}_3(1) = \mathcal{E}_3(1).$$

On the other hand denote by $\text{int}_{A_c}(\mathcal{E}'_3 \cap A_c)$ the interior of the set $\mathcal{E}'_3 \cap A_c$ (subspace topology) for $|c| < 1$, *i.e.*, the open triangle defined in Lemma 2.4, and observe that

$$\text{int } \mathcal{E}'_3 = \bigcup_{|c| < 1} \text{int}_{A_c}(\mathcal{E}'_3 \cap A_c)$$

as we can find a neighborhood around each point of $\text{int}_{A_c}(\mathcal{E}'_3 \cap A_c)$, $|c| < 1$ which is contained in \mathcal{E}'_3 . Further each point of $\text{int}_{A_c}(\mathcal{E}'_3 \cap A_c)$ satisfies the conditions of $\mathcal{E}_3(1)$ whenever $|c| < 1$. Hence

$$\text{int } \mathcal{E}'_3 = \bigcup_{|c| < 1} \text{int}(\mathcal{E}'_3 \cap A_c) \subset \mathcal{E}_3(1).$$

Thus we have shown that $\text{int } \mathcal{E}'_3 = \mathcal{E}_3(1)$.

To prove the theorem we show $\mathcal{E}'_3 = \overline{\text{int } \mathcal{E}'_3}$. We already have that $\text{int } \mathcal{E}'_3 = \bigcup_{|c| < 1} \text{int}_{A_c}(\mathcal{E}'_3 \cap A_c)$. Hence we look at the convergent sequences of points out of $\bigcup_{|c| < 1} \text{int}(\mathcal{E}'_3 \cap A_c)$. Such a sequence converges either to some point within $\bigcup_{|c| < 1} (\mathcal{E}'_3 \cap A_c)$ or to some point within one of the sets $\lim_{c \rightarrow \pm 1} (\mathcal{E}'_3 \cap A_c)$. From Lemma 2.4 we already have

$$\mathcal{E}'_3 \cap A_c = \Delta((c, -1, -c)(c, 1 - 2c, c - 2), (c, 2c + 1, c + 2))$$

and we see that

$$\begin{aligned} \lim_{c \rightarrow 1} (\mathcal{E}'_3 \cap A_c) &= \{(1, \lambda, \lambda) \mid -1 \leq \lambda \leq 3\}, \\ \lim_{c \rightarrow -1} (\mathcal{E}'_3 \cap A_c) &= \{(-1, \lambda, -\lambda) \mid -1 \leq \lambda \leq 3\} \end{aligned}$$

which exactly correspond to the sets $(\mathcal{E}'_3 \cap A_{\pm 1})$. Thus

$$\overline{\mathcal{E}_3(1)} = \overline{\text{int } \mathcal{E}'_3} = \bigcup_{|c| \leq 1} (\mathcal{E}'_3 \cap A_c) = \mathcal{E}'_3$$

and we are done. \square

Finally we have a representation of the closure of $\mathcal{E}_3(1)$. In the proof of Lemma 2.4 we already recognized that the number of inequalities to describe \mathcal{E}'_3 can be reduced. Indeed, by using an algorithm (Algorithm 3) which we will present in Section 4, we gain

$$\overline{\mathcal{E}_3(1)} = \{(x, y, z) \mid |x + z| \leq 1 + y \wedge y - xz \leq 1 - x^2 \wedge |z| \leq 3\}.$$

3. STATEMENT OF THE MAIN RESULT

In this section we give a complete description of \mathcal{D}_3^0 . To this matter we define the sets

$$S_1 := \{(x, y, z) \mid 2x - 2z \geq 1 \wedge 2x + 2y + 2z > -1 \wedge 2x + 2y \leq 1 \wedge 2x \leq 1 \wedge 2x - 2y + 2z \leq 1\},$$

$$S_2 := \{(x, y, z) \mid x - z \leq -1 \wedge 2x - 2y + 2z \leq 1 \wedge -2x + 2y \leq 1 \wedge 2x > -1\},$$

$$S_3 := \{(x, y, z) \mid x - z > -1 \wedge 2x - 2y + 2z \leq 1 \wedge -2x + 2y < 1, 2x > -1 \wedge 2x - 2z < -1 \\ \wedge 2x + 2y + 2z > -1\},$$

$$S_4 := \{(x, y, z) \mid 2x - 2y + 2z \leq 1 \wedge -2x + 2y \leq 1 \wedge 2x - 2z = -1 \wedge 2x + 2y + 2z > -1\},$$


$$S_5 := \{(x, y, z) \mid -1 < 2x \leq 1 \wedge -1 < 2x - 2z \leq 1 \wedge 2x + 2y + 2z > -1 \wedge 2x - 2y + 2z \leq 1 \\ \wedge 2x + 4y - 2z < 3, 2y \leq 1\}$$

and denote their union by

$$\mathcal{S} := \bigcup_{i \in \{1, \dots, 5\}} S_i.$$

Note that S_1, S_2, S_3, S_5 are polyhedra while S_4 is a polygon. The following theorem states the main result of the present paper.

Theorem 3.1. $\mathcal{D}_3^0 = \mathcal{S}$

FIGURE 4. A view of \mathcal{D}_3^0

Two views of the set \mathcal{D}_3^0 are depicted in Figure 4 and Figure 5. For rotating 3D-pictures of \mathcal{D}_3^0 we refer the reader to the authors' home pages [16].

In Section 4 we will prove this theorem. Here we want to give an outline of the proof. In a first step we will use Theorem 2.2 in order to show that

$$(3.1) \quad \mathcal{S} \subseteq \mathcal{D}_3^0.$$

For showing the opposite inclusion we need a set of nonzero periods Π such that for $\mathcal{P} := \bigcup_{\pi \in \Pi} P(\pi)$ we have

$$\mathcal{S} \cup \mathcal{P} \supseteq \mathcal{D}_3.$$

From (3.1) we can deduce $\mathcal{S} \cap \mathcal{P} = \emptyset$. Thus,

$$\mathcal{S} \supseteq \mathcal{D}_3 \setminus \mathcal{P} \supseteq \mathcal{D}_3^0.$$


Since $\mathcal{D}_3 \subset \overline{\mathcal{E}_3(1)}$ we are done if we can cover $\overline{\mathcal{E}_3(1)}$ with $\mathcal{P} \cup \mathcal{S}$, *i.e.*, if we can show that

$$\mathcal{P} \cup \mathcal{S} \supseteq \overline{\mathcal{E}_3(1)}.$$

4. PROOF OF THE MAIN RESULT

We will prove our result in two parts according to the outline given in the previous section. First of all, we set up some notations.

Notation 4.1. For a logical system \mathcal{J} of inequalities, which are combined by \wedge and \vee , denote by $X(\mathcal{J})$ the set of all points that satisfy \mathcal{J} . Let P a set of inequalities. Then $\bigwedge P$ and $\bigvee P$ denote the systems $\bigwedge_{I \in P} I$ and $\bigvee_{I \in P} I$, respectively.

FIGURE 5. A view of \mathcal{D}_3^0

For the rest of the section denote by T_i the set of inequalities that define the set S_i for $i \in \{1, \dots, 5\}$. These sets are assembled only of single inequalities. We have

$$\begin{aligned}
 T_1 &:= \{2x - 2z \geq 1, 2x + 2y + 2z > -1, 2x + 2y \leq 1, 2x \leq 1, 2x - 2y + 2z \leq 1\}, \\
 T_2 &:= \{x - z \leq -1, 2x - 2y + 2z \leq 1, -2x + 2y \leq 1, 2x > -1\}, \\
 T_3 &:= \{x - z > -1, 2x - 2y + 2z \leq 1, -2x + 2y < 1, 2x > -1, 2x - 2z < -1, 2x + 2y + 2z > -1\}, \\
 T_4 &:= \{2x - 2y + 2z \leq 1, -2x + 2y \leq 1, 2x - 2z \leq -1, 2x - 2z \geq -1, 2x + 2y + 2z > -1\}, \\
 T_5 &:= \{-1 < 2x, 2x \leq 1, -1 < 2x - 2z, 2x - 2z \leq 1, 2x + 2y + 2z > -1, \\
 &\quad 2x - 2y + 2z \leq 1, 2x + 4y - 2z < 3, 2y \leq 1\},
 \end{aligned}$$

hence the equality of S_4 and the two double inequalities of S_5 are split into inequalities. Thus, $S_i = X(\bigwedge T_i)$ for $i = 1, \dots, 5$. Denote by \bar{T}_i the set T_i with all the strict inequalities changed to not strict ones. Since all occurring inequalities are linear it can easily be checked that $\bar{S}_i = X(\bigwedge \bar{T}_i)$.

Table 1 shows 43 different periods, we denote the corresponding polyhedron by $P(\pi_j)$, where $j \in \{1, \dots, 43\}$.

Now for each $i \in \{1, \dots, 43\}$ define Q_i to be the set of single inequalities such that $P(\pi_i) = X(\bigwedge Q_i)$. For instance, the set Q_{19} can be defined by

$$\begin{aligned}
 Q_{19} &:= \left\{ -\frac{1}{2} \leq x + y - z - 1, x + y - z - 1 < \frac{1}{2}, -\frac{1}{2} \leq x - y - z, x - y - z < \frac{1}{2}, \right. \\
 &\quad \left. -\frac{1}{2} \leq -x - y + 1, -x - y + 1 < \frac{1}{2}, -\frac{1}{2} \leq -x + z + 1, \right. \\
 &\quad \left. -x + z + 1 < \frac{1}{2}, -\frac{1}{2} \leq y + z - 1, y + z - 1 < \frac{1}{2} \right\}
 \end{aligned}$$

Length	Periods			
3	$\pi_1=(-1, -1, -1)$ $\pi_5=(0, -1, 1)$	$\pi_2=(-1, -1, 0)$	$\pi_3=(-1, 0, 1)$	$\pi_4=(0, -1, 0)$
4	$\pi_6=(0, -1, 0); -1$	$\pi_7=(0, -1, 0); 1$	$\pi_8=(1, -1, 1); -1$	
5	$\pi_9=(-2, 1, -1); -1, 1$	$\pi_{10}=(-2, 1, 0); -1, 2$	$\pi_{11}=(-1, -1, 1); 1, 0$	
	$\pi_{12}=(0, -2, -1); 1, 2$	$\pi_{13}=(0, -1, 1); -1, 0$	$\pi_{14}=(0, 1, -1); 1, 0$	
	$\pi_{15}=(0, 1, 0); -1, -1$	$\pi_{16}=(0, 1, 0); -1, 0$	$\pi_{17}=(0, 2, 1); -1, -2$	
	$\pi_{18}=(1, -1, 1); -1, 0$	$\pi_{19}=(1, 1, -1); -1, 0$	$\pi_{20}=(2, -1, 0); 1, -2$	
6	$\pi_{21}=(0, -1, 0); 0, 1, 0$		$\pi_{22}=(1, 1, 0); -1, -1, 0$	
7	$\pi_{23}=(0, 1, -1); -1, 1, 0, -1$		$\pi_{24}=(1, 1, 0); -1, -1, -1, 0$	
8	$\pi_{25}=(-1, -1, 1); 1, 2, 0, 0, -2$		$\pi_{26}=(-1, 0, 0); 1, 0, 0, -1, -1$	
	$\pi_{27}=(-1, 1, 0); -1, 1, -1, 0, 1$		$\pi_{28}=(0, 0, 2); 1, 1, -1, -1, -2$	
	$\pi_{29}=(1, 1, 1); 0, -1, -1, -1, 0$		$\pi_{30}=(2, 1, -1); -2, -2, -1, 1, 2$	
9	$\pi_{31}=(-1, 0, 0); 1, 1, 1, 0, -1, -1$		$\pi_{32}=(0, 1, 1); 1, 0, -1, -2, -2, -1$	
10	$\pi_{33}=(-1, -1, 1); 0, -1, 1, 1, -1, 0, 1$		$\pi_{34}=(0, -2, 1); 1, -2, 0, 2, -1, -1, 2$	
	$\pi_{35}=(0, -1, -1); -1, 0, 0, 1, 1, 1, 0$		$\pi_{36}=(1, 2, 1); 1, -1, -1, -2, -1, -1, 1$	
	$\pi_{37}=(1, 2, 2); 1, 0, -1, -2, -2, -1, 0$			
11	$\pi_{38}=(-2, 0, 1); -2, 1, 0, -2, 2, -1, -1, 2$			
	$\pi_{39}=(0, 1, 2); 2, 1, 0, -1, -2, -2, -2, -1$			
12	$\pi_{40}=(-2, 2, -1); 0, 1, -2, 2, -2, 1, 0, -1, 2$			
	$\pi_{41}=(0, 1, 2); 2, 2, 1, 0, -1, -2, -2, -2, -1$			
13	$\pi_{42}=(0, 1, -2); 2, -1, -1, 2, -2, 1, 0, -1, 1, -1$			
22	$\pi_{43}=(0, 2, 2); 1, -1, -2, -2, 0, 1, 2, 1, 0, -2, -2, -1, 1, 2, 2, 0, -1, -2, -1$			

TABLE 1. The 43 periods needed to cutout \mathcal{D}_3^0

(see also Example 2.3). Finally we set

$$\mathcal{P} := \bigcup_{j=1}^{43} P(\pi_j).$$

Remark 4.2. We note that the construction of the set \mathcal{S} as well as the exhibition of the 43 periods corresponding to relevant cutout polyhedra has been achieved by extensive computer experiments. Up to now we do not know an easy way that would lead to a list of all the cutouts needed to get the set \mathcal{D}_3^0 . To find an algorithmic way to construct all these cutouts is desirable since it could lead to characterizations of \mathcal{D}_d^0 even for higher dimensions d .

Observe that no element of the 43 periods given above contains elements having modulus greater than 2. Up to now, we do not know the reason for this fact. In order to characterize $\tilde{\mathcal{D}}_2^0$ we need periods with elements that are arbitrarily large (cf. [1, Sections 6 and 7]).

4.1. Using the algorithm of Section 2. Theorem 2.2 shows the existence of an algorithm for the construction of a graph $G(H) = V \times E$ which can be used for finding all periods within the convex body H . Following [5], the graph is constructed recursively. Define $H = \square(\mathbf{r}_1, \dots, \mathbf{r}_k) \subset \text{int } \mathcal{D}_3$ to be the convex hull of some points $\mathbf{r}_1, \dots, \mathbf{r}_k$. For a $\mathbf{z} \in \mathbb{Z}^d$, let $m(\mathbf{z}) = \min_{i \in \{1, \dots, k\}} (-\lfloor \mathbf{r}_i \mathbf{z} \rfloor)$ and $M(\mathbf{z}) = \max_{i \in \{1, \dots, k\}} (-\lfloor \mathbf{r}_i \mathbf{z} \rfloor)$. Set

$$V_0 := \{\pm \mathbf{e}_i \mid i = 1, \dots, d\}$$

and then successively calculate V_1, V_2, \dots by the rule

$$V_{i+1} := V_i \cup \{(z_2, \dots, z_d, j) \mid \mathbf{z} = (z_1, \dots, z_d) \in V_i, -M(-\mathbf{z}) \leq j \leq M(\mathbf{z})\}.$$

For sets H having a sufficiently small diameter the iteration stabilizes yielding $V := V_n = V_{n+1}$ for some $n \in \mathbb{N}$. The set of edges is constructed by

$$E := \{(\mathbf{x}, (z_2, \dots, z_d, j)) \mid \mathbf{x} = (z_1, \dots, z_d) \in V, m(\mathbf{z}) \leq j \leq M(\mathbf{z})\}.$$

Let \mathcal{Q} be a system of linear, non-strict inequalities linked with \wedge . Then $X(\mathcal{Q})$ forms a convex polyhedron that can be regarded as the convex hull of finitely many points $\mathbf{r}_1, \dots, \mathbf{r}_k$. Suppose that $X(\mathcal{Q}) \subset \mathcal{E}_3(1)$. We want to set up an algorithm that calculates the set of all periods π whose associated polyhedron $P(\pi)$ has non-empty intersection with $X(\mathcal{Q})$. Theorem 2.2 ensures the existence of such an algorithm only if $X(\mathcal{Q})$ has sufficiently small diameter. If the set $X(\mathcal{Q})$ is too big, the graph $G(X(\mathcal{Q}))$ is infinite. We solve this problem in the following way. Suppose that, during the calculation of $|V|$, we obtain a set V_i whose number of elements $|V_i|$ exceeds an appropriate bound p . In this case we stop the calculation of V and divide the set $X(\mathcal{Q})$ into two parts for which we calculate the set V again. By recursively doing this splitting procedure we eventually end up with sets whose diameter is small enough (provided that p is chosen reasonably).

Suppose that the set $X(\mathcal{Q})$ is the convex hull of its k vertices $\mathbf{r}_1, \dots, \mathbf{r}_k$. We do not know these vertices explicitly. What we need is just $m(\mathbf{z})$ and $M(\mathbf{z})$ for certain fixed values of $\mathbf{z} \in \mathbb{Z}^d$. However, as \mathcal{Q} is given as a system of linear inequalities, we easily see that

$$\begin{aligned} m(z) &= \min_{\mathbf{r} \in X(\mathcal{Q})} (-\lfloor \mathbf{r}\mathbf{z} \rfloor), \\ M(z) &= \max_{\mathbf{r} \in X(\mathcal{Q})} (-\lfloor \mathbf{r}\mathbf{z} \rfloor). \end{aligned}$$

The extremal values on the left hand side can now easily be calculated by standard linear optimization.

The algorithm consists of two parts. The first part is Algorithm 1, which constructs the set of vertices V of the graph $G(X(\mathcal{Q}))$ for a given convex body $X(\mathcal{Q})$. Whenever during the calculation the size of this set exceeds a given bound p , Algorithm 1 stops returning an overflow. Otherwise it terminates by returning V . Denote the application of Algorithm 1 with parameter \mathcal{Q} and bound p by $\text{VG}(\mathcal{Q}, p)$ (VG = **v**ertices of the **g**raph). Algorithm 2 is recursive. As input we

Algorithm 1 Calculation of the set of vertices of $G(X(\mathcal{Q}))$: VG

Input: \mathcal{Q}, p

Output: set of vertices V

```

1:  $V \leftarrow \{\pm \mathbf{e}_j | j = 1, \dots, d\}$ 
2:  $M \leftarrow \emptyset$ 
3: while  $V \neq M$  do
4:   if  $\#V > p$  then
5:     return(Overflow)
6:     stop calculation
7:   end if
8:    $N \leftarrow V \setminus M$ 
9:    $M \leftarrow V$ 
10:  for all  $(x_1, \dots, x_d) \in N$  do
11:     $i \leftarrow \min_{(r_1, \dots, r_d) \in X(\mathcal{Q})} (\lfloor -\sum_{k=1}^d x_k r_k \rfloor)$ 
12:     $j \leftarrow \max_{(r_1, \dots, r_d) \in X(\mathcal{Q})} (-\lfloor \sum_{k=1}^d x_k r_k \rfloor)$ 
13:     $V \leftarrow V \cup \{(x_2, \dots, x_d, k) | k \in \{i, \dots, j\}\}$ 
14:  end for
15: end while
16: return( $V$ )

```

have \mathcal{Q} and we write $\text{FP}(\mathcal{Q})$ for its application on \mathcal{Q} (FP = **f**ind all **p**eriods). Algorithm 2 evokes Algorithm 1 to calculate the set of vertices of $G(X(\mathcal{Q}))$. If an overflow occurs, the set $X(\mathcal{Q})$ is split with respect to some hyperplane $G(X_1, \dots, X_d) = 0$. Then Algorithm 2 is applied on $\mathcal{Q}_1 := (\mathcal{Q} \wedge G(X_1, \dots, X_d) \leq 0)$ and $\mathcal{Q}_2 := (\mathcal{Q} \wedge G(X_1, \dots, X_d) \geq 0)$ separately. If there is no overflow and V is returned, the set of edges E is calculated and all the cycles are extracted. These cycles induce the periods, we are searching for. Note that the subsets \mathcal{Q}_1 and \mathcal{Q}_2 are again defined by finitely many non-strict inequalities so that they can be treated by Algorithm 1 in the same way as \mathcal{Q} .

Algorithm 2 Search for all periods within an area $X(\mathcal{Q})$ (recursively): FP

Input: \mathcal{Q}

Output: Π list of cycles

```

1:  $p \leftarrow$  suitable bound
2:  $V \leftarrow \text{VG}(\mathcal{Q}, p)$ 
3: if  $\neg(\text{overflow})$  then
4:    $E \leftarrow$  set of edges of  $G(X(\mathcal{Q}))$ 
5:    $\Pi \leftarrow$  periods induced by the cycles of  $G(X(\mathcal{Q}))$ 
6: else
7:   construct  $\mathcal{Q}_1, \mathcal{Q}_2$ 
8:    $\Pi \leftarrow \text{FP}(\mathcal{Q}_1)$ 
9:    $\Pi \leftarrow \Pi \cup \text{FP}(\mathcal{Q}_2)$ 
10: end if
11: return( $\Pi$ )

```

In our setting we need to apply Algorithm 2 to the sets defined by the inequalities \bar{T}_i ($i \in \{1, \dots, 5\}$). All we need to specify is the subdividing strategy and the bound p for $|V|$. As for the subdividing strategy we subdivide a given set in two parts by a plane orthogonal to the coordinate axis in which the set has its biggest extension. To make this precise, let

$$m_i := \min_{(x_1, x_2, x_3) \in X(\mathcal{Q})} x_i, i = 1, 2, 3,$$

$$M_i := \max_{(x_1, x_2, x_3) \in X(\mathcal{Q})} x_i, i = 1, 2, 3,$$

and $j \in \{1, 2, 3\}$ be the smallest index for which $M_j - m_j = \max(M_1 - m_1, M_2 - m_2, M_3 - m_3)$. The dividing hyperplane is now defined by $G(X_1, X_2, X_3) = 0$ with $G(X_1, X_2, X_3) := X_j - \frac{M_j + m_j}{2}$.

For the upper bound of the number of vertices it turns out that a choice depending on the quantities $M_j - m_j$ is convenient. In particular, we choose $p = \frac{20}{M_j - m_j}$. Then we get the following result

Lemma 4.3. $\text{FP}(\bigwedge T_i)$ terminates for each $i \in \{1, \dots, 5\}$.

Proof. We implemented the algorithms for T_i with the above mentioned subdivision strategy and bounds in Mathematica[®]. The program is available on the authors' homepages [16]. \square

Theorem 4.4. $S_i \subset \mathcal{D}_3^0$ holds for all $i \in \{1, \dots, 5\}$.

Proof. For each $i \in \{1, \dots, 5\}$ we have that $X(\bigwedge \bar{T}_i)$ is a convex hull of finitely many points. Moreover, $X(\bigwedge \bar{T}_i) = \bar{S}_i$. Denote by Π_i the set of periods computed by the application of Algorithm 2 on $\bigwedge \bar{T}_i$. Hence Π_i includes all periods associated to polyhedra having non-empty intersection with $X(\bigwedge \bar{T}_i)$. Now, according to (2.2), each of these periods $\pi \in \Pi_i$ induces a system of inequalities $\mathcal{P}(\pi)$. It turns out that for each $\pi \in \Pi_i$ we have

$$X(\mathcal{P}(\pi) \wedge \bigwedge T_i) = \emptyset \text{ holds for each } i \in \{1, \dots, 5\}$$

(an easy way for checking this is to apply the cylindrical algebraic decomposition algorithm). Thus there is no period that yields a nonempty cutout intersecting with S_i and therefore $S_i \subset \mathcal{D}_3^0$. \square

4.2. Covering the set $\mathcal{D}_3 \setminus \mathcal{D}_3^0$ by cutout polyhedra. Fix Q_1, \dots, Q_{43} to be the sets of inequalities of the 43 polyhedra induced by the periods given in Table 1, where Q_j denotes just the reduced set of inequalities such that $X(\bigwedge Q_j)$ yields the corresponding polyhedron for any $j \in \{1, \dots, 43\}$. "Reduced" means that all the redundant inequalities are removed.

Remark 4.5. It is not really necessary to work with the reduced systems but the main algorithm works much faster and the reduction is not too difficult to realize.

The algorithm simply uses the fact that an inequality I is redundant for a system $\mathcal{S} \wedge I$ if $X(\mathcal{S} \wedge I) = X(\mathcal{S})$ or, equivalently, $X(\mathcal{S} \wedge \neg I) = \emptyset$. Denote the application of Algorithm 3 with parameter P by $\text{RL}(P)$ (RL=reduce list of inequalities).

Algorithm 3 Reducing a list of inequalities: RL

Input: P set of inequalities**Output:** P reduced set of inequalities

```

1: for all inequalities  $I \in P$  do
2:    $P \leftarrow P \setminus I$ 
3:   if  $X(\bigwedge P \wedge \neg I) \neq \emptyset$  then
4:      $P \leftarrow P \cup I$ 
5:   end if
6: end for
7: return( $P$ )

```

At the end of Section 2 we found a parametrization of $\overline{\mathcal{E}_3(1)}$. We saw that $\overline{\mathcal{E}_3(1)} = X(\bigwedge D)$ for

$$D := \{x + z \leq 1 + y, -1 - y \leq x + z, y - xz \leq 1 - x^2, z \leq 3, z \geq -3\}.$$

Let \mathcal{P} be a list of sets of inequalities and G to be a set of inequalities. We want to verify if $\bigcup_{P \in \mathcal{P}} X(\bigwedge P)$ covers $X(\bigwedge G)$. This is equivalent to

$$(4.1) \quad X\left(\bigwedge G \wedge \neg \bigvee_{P \in \mathcal{P}} \bigwedge P\right) = \emptyset.$$

In principle we could do this verification directly. For computational reasons we are a little more restricted. (In fact the direct verification of (4.1) overcharges Mathematica[®]). A verification of a claim of the shape (4.1) can be done in a reasonable amount of time if $\#\mathcal{P} \leq 3$. We give an algorithm that solves this problem for general \mathcal{P} and G by a subdivision process. The idea is to split the set $X(\bigwedge G)$ into suitable subsets and hope that each of these subsets is covered by a smaller number of sets. First we state Algorithm 4 which removes those sets from \mathcal{P} that do not affect G , hence a set P is removed when $X(\bigwedge G) \cap X(\bigwedge P) = \emptyset$. Denote the application of this algorithm by $\text{RS}(G, \mathcal{P})$ (RS=remove inequalities with respect to a set).

Algorithm 4 Removing those lists of inequalities that do not affect a given set G : RS

Input: G, \mathcal{P} **Output:** \mathcal{P} reduced list of inequalities

```

1: for all sets  $P \in \mathcal{P}$  do
2:   if  $X(\bigwedge G \wedge \bigwedge P) = \emptyset$  then
3:      $\mathcal{P} \leftarrow \mathcal{P} \setminus P$ 
4:   end if
5: end for
6: return( $\mathcal{P}$ )

```

The main algorithm (Algorithm 5) is recursive. As an input we have again \mathcal{P} and G of the usual shape, where \mathcal{P} is reduced by Algorithm 4. Whenever the algorithm recognizes that a subset of $X(\bigwedge G)$ is not fully covered by the sets described in \mathcal{P} , it returns this subset. Denote the application by $\text{VC}(G, \mathcal{P})$ (VC=verify covering). At first Algorithm 5 checks whether $\#\mathcal{P} \leq 3$. If this is the case we can verify whether (4.1) holds, otherwise we choose an arbitrary inequality $I \in \bigcup_{P \in \mathcal{P}} P$ such that $X(\bigwedge G \wedge I) \neq X(\bigwedge G)$. There are two possibilities:

- There is such an inequality I . Then $X(\bigwedge G)$ is split by adding I and $\neg I$, respectively, to G and Algorithm 5 is applied (recursively) on both of these subsets. Algorithm 4 is used to possibly reduce \mathcal{P} for each of the subsets. These reduced sets form the second parameter.
- There is no such I . But this would mean that all the points of $X(\bigwedge G)$ suffice all inequalities of $\bigcup_{P \in \mathcal{P}} P$. This is equivalent to $X(\bigwedge G) \subset X(P)$ for any $P \in \mathcal{P}$ and this implies that G and \mathcal{P} suffice the condition (4.1).

Now, whenever (4.1) is not fulfilled, the set $X(\wedge G)$ is not covered by $X(\bigvee_{P \in \mathcal{P}} \wedge P)$ and the algorithm returns the affected set $X(\wedge G)$. The application of Algorithm 5 terminates without any output if $X(\bigvee_{P \in \mathcal{P}} \wedge P)$ covers $X(\wedge G)$.

Algorithm 5 Checks if a set is covered by the union of others (recursively): VC

Input: G, \mathcal{P}

Output: subsets of $X(\wedge G)$ that are not fully covered by $X(\bigvee_{P \in \mathcal{P}} \wedge P)$

```

1: if  $\#\mathcal{P} \leq 3$  then
2:   if  $X(G \wedge \neg \bigvee_{P \in \mathcal{P}} \wedge P) \neq \emptyset$  then
3:     return  $X(\wedge G)$  is not fully covered
4:   end if
5: else
6:   if  $\exists I \in \bigcup_{P \in \mathcal{P}} P : X(\wedge G \wedge I) \neq \emptyset$  then
7:     VC(RL( $G \cap \{I\}$ ), RS( $G \cap \{I\}$ ),  $\mathcal{P}$ )
8:     VC(RL( $G \cap \{\neg I\}$ ), RS( $G \cap \{\neg I\}$ ),  $\mathcal{P}$ )
9:   end if
10: end if

```

We can now state the main theorem of this subsection.

Theorem 4.6. *The algorithm VC(D, \mathcal{P}) terminates without yielding any output for*

$$\mathcal{P} = \{Q_1, \dots, Q_{43}, T_1, \dots, T_5\}.$$

Proof. We implemented the algorithms in Mathematica[®]. The program is available on the authors' homepages [16]. \square

Theorem 4.6 shows all the periods together with our set to really cover all of $\overline{\mathcal{E}_3(1)}$ and thus cover \mathcal{D}_3 . Thus, the cutout polyhedra $P(\pi_1), \dots, P(\pi_{43})$ cover the whole set $\overline{\mathcal{E}_3(1)} \setminus \mathcal{S}$. Hence, in view of Theorem 4.4 we get that

$$\overline{\mathcal{E}_3(1)} \setminus \mathcal{S} \subset \bigcup_{1 \leq i \leq 43} P(\pi_i).$$

Together with Theorem 4.4 this yields Theorem 3.1 and we are done.

REFERENCES

- [1] S. AKIYAMA, T. BORBÉLY, H. BRUNOTTE, A. PETHŐ AND J. M. THUSWALDNER, *Generalized radix representations and dynamical systems I*, Acta Math. Hungar. 108 (2005), 207–238.
- [2] S. AKIYAMA, H. BRUNOTTE, A. PETHŐ AND W. STEINER, *Remarks on a conjecture on certain integer sequences*, Period. Math. Hungar. 52, Nr. 1 (2006), 1–17.
- [3] S. AKIYAMA, H. BRUNOTTE, A. PETHŐ AND J. M. THUSWALDNER, *Generalized radix representations and dynamical systems II*, Acta Arithmetica 121 (2006), 21–61.
- [4] S. AKIYAMA, H. BRUNOTTE, A. PETHŐ AND J. M. THUSWALDNER, *Generalized radix representations and dynamical systems III*, preprint.
- [5] S. AKIYAMA, K. SCHEICHER, *Symmetric shift radix systems and finite expansions*, preprint.
- [6] H. BRUNOTTE, *On trinomial bases of radix representations of algebraic integers*, Acta Sci. Math. Acta Sci. Math. (Szeged) 67 (2001), 521–527.
- [7] G. E. COLLINS *Quantifier Elimination for the Elementary Theory of Real Closed Fields by Cylindrical Algebraic Decomposition*. Lect. Notes Comput. Sci. 33 (1975) 134–183.
- [8] C. FROUGNY, B. SOLOMYAK, *Finite beta-expansions*, Ergod. Th. and Dynam. Sys. 12 (1992), 713–723.
- [9] T. S. MOTZKIN, H. RAIFFA, G. L. THOMPSON, R. L. THRALL, *The double description method*. Contributions to the theory of games, vol. 2, pp. 51–73. Annals of Mathematics Studies, no. 28. Princeton University Press, Princeton, N. J., (1953).
- [10] W. PARRY, *On the β -expansions of real numbers*, Acta Math. Acad. Sci. Hungar. 11 (1960), 401–416.
- [11] S. PEMMARAJU, S. SKIENA, *Computational Discrete Mathematics, Combinatorics and Graph Theory with Mathematica[®]*, Cambridge University Press (2003).
- [12] A. PETHŐ, *On a polynomial transformation and its application to the construction of a public key cryptosystem*, in: Computational Number Theory, A. Pethő, M. Pohst, H. C. Williams and H. G. Zimmer (eds.), de Gruyter (1991), 31–43.

- [13] A. RÉNYI, *Representations for real numbers and their ergodic properties*, Acta Math. Acad. Sci. Hungar. 8 (1957), 477–493.
- [14] I. SCHUR, *Über Potenzreihen, die im inneren des Einheitskreises beschränkt sind*, J. reine angew. Math. 148 (1918), 122–145.
- [15] P. SURER, *New characterisation results for shift radix systems*, preprint.
- [16] P. SURER, Shift Radix Systems, <http://www.unileoben.ac.at/~mathstat/personal/surer.htm>.
- [17] T. TAKAGI, *Lectures in Algebra*, (1965).
- [18] R. TARJAN, *Depth-first search and linear graph algorithms*, SIAM J. Comput. 1 (1972), no. 2, 146–160.

FACULTY OF INFORMATICS, UNIVERSITY OF DEBRECEN, P.O. BOX 12, H-4010 DEBRECEN, HUNGARY
E-mail address: `husztia@inf.unideb.hu`

JOHANN RADON INSTITUTE FOR COMPUTATIONAL AND APPLIED MATHEMATICS (RICAM), AUSTRIAN ACADEMY OF SCIENCES, ALTENBERGERSTRASSE 69, A-4040 LINZ, AUSTRIA
E-mail address: `klaus.scheicher@oeaw.ac.at`

CHAIR OF MATHEMATICS AND STATISTICS, FRANZ-JOSEF-STRASSE 18, UNIVERSITY OF LEOBEN, A-8700 LEOBEN, AUSTRIA
E-mail address: `paul.surer@reflex.at`

CHAIR OF MATHEMATICS AND STATISTICS, UNIVERSITY OF LEOBEN, FRANZ-JOSEF-STRASSE 18, A-8700 LEOBEN, AUSTRIA
E-mail address: `joerg.thuswaldner@unileoben.ac.at`