

Critical yield numbers and limiting yield surfaces of particle arrays settling in a Bingham fluid

J.A. Iglesias, G. Mercier, O. Scherzer

RICAM-Report 2018-05

Critical yield numbers and limiting yield surfaces of particle arrays settling in a Bingham fluid

José A. Iglesias ^{*1}, Gwenael Mercier ^{†1}, and Otmar Scherzer ^{‡1,2}

¹Johann Radon Institute for Computational and Applied Mathematics (RICAM)
Austrian Academy of Sciences, Linz, Austria.

²Computational Science Center, University of Vienna, Vienna, Austria.

Abstract

We consider the flow of multiple particles in a Bingham fluid in an anti-plane shear flow configuration. The limiting situation in which the internal and applied forces balance and the fluid and particles stop flowing, that is, when the flow settles, is formulated as finding the optimal ratio between the total variation functional and a linear functional. The minimal value for this quotient is referred to as the critical yield number or, in analogy to Rayleigh quotients, generalized eigenvalue. This minimum value can in general only be attained by discontinuous, hence not physical, velocities. However, we prove that these generalized eigenfunctions, whose jumps we refer to as limiting yield surfaces, appear as rescaled limits of the physical velocities. Then, we show the existence of geometrically simple minimizers. Furthermore, a numerical method for the minimization is then considered. It is based on a nonlinear finite difference discretization, whose consistency is proven, and a standard primal-dual descent scheme. Finally, numerical examples show a variety of geometric solutions exhibiting the properties discussed in the theoretical sections.

1 Introduction

In this article, we investigate the stationary flow of particles in a Bingham fluid. Such fluids are important examples of non-Newtonian fluids, describing for instance cement, toothpaste, and crude oil [24]. They are characterized by two numerical quantities: a yield stress τ_Y that must be exceeded for strain to appear, and a fluid viscosity μ_f that describes its linear behaviour once it starts to flow (see figure 1).

An important property of Bingham fluid flows is the occurrence of plugs, which are regions where the fluid moves like a rigid body. Such rigid movements occur at positions where the stress does not exceed the yield stress.

In this paper we consider anti-plane shear flow in an infinite cylinder, where an ensemble of inclusions move under their own weight inside a Bingham fluid of lower density, and in which the gravity and viscous forces are in equilibrium (cf (5)), therefore inducing a flow which is *steady* or *stationary*, that is, in which the velocity does not depend on time. For such a configuration, we are interested in determining the ratio between applied forces and the yield stress such that the Bingham fluid stops flowing completely. This ratio is called *critical yield number*.

*jose.iglesias@ricam.oeaw.ac.at

†gwenael.mercier@ricam.oeaw.ac.at

‡otmar.scherzer@univie.ac.at




Figure 1: Relation between stress and strain in a Bingham fluid

Related work.

To our knowledge, the first mathematical studies of critical yield numbers were conducted by Mosolov & Miasnikov [20, 21], who also considered the anti-plane situation for flows inside a pipe. In particular, they discovered the geometrical nature of the problem and related the critical yield number to what in modern terminology is known as the Cheeger constant of the cross-section of the region containing the fluid. Very similar situations appear in the modelling of the onset of landslides [13, 15, 12], where non-homogeneous coefficients and different boundary conditions arise. Two-fluid anti-plane shear flows that arise in oilfield cementing are studied in [10, 11]. Settling of particles under gravity, not necessarily in anti-plane configurations is also considered in [16, 23]. Finally, the previous work [9] also focuses in the anti-plane settling problem. There, the analysis is limited to the case in which all particles move with the same velocity and where the main interest is to extract the critical yield numbers from geometric quantities. In the current work we lift this restriction and focus on the calculations of the limiting velocities, also from a numerical point of view. Various applications of the critical yield stress of suspensions are pointed out in [4, Section 4.3].

Structure of the paper.

We begin in Section 2 by recalling the mathematical models describing the stationary Bingham fluid flow in an anti-plane configuration, and an optimization formulation for determining the critical yield number.

Next, in Section 3 we consider a relaxed formulation of this optimization problem, which is naturally set in spaces of functions of bounded variation, and show that the limiting velocity profile as the flow stops is a minimizer of this relaxed problem.

In Section 4, as in the case of a single particle [9], we prove that there exists a minimizer that attains only two non zero velocity values.

Finally, in Section 5 we present a numerical approach to compute minimizers. This approach is based on the non-smooth convex optimization scheme of Chambolle-Pock [8] and an upwind finite difference discretization [7]. We prove the convergence of the discrete minimizers to continuous ones as the grid size decreases to zero. We then use this scheme to illustrate the theoretical results of Section 4.

2 The model

We consider a Bingham fluid filling a vertical cylindrical domain $\hat{\Omega} \times \mathbb{R} \subset \mathbb{R}^3$ and a solid inclusion $\hat{\Omega}_s \times \mathbb{R} \subset \hat{\Omega} \times \mathbb{R}$. We denote by $\hat{\Omega}_f = \hat{\Omega} \setminus \hat{\Omega}_s$ the portion of the domain occupied by the fluid, and by ρ_s, ρ_f the corresponding constant densities. We focus on a vertical stationary flow, meaning that the solid flows down at a vertical velocity $\hat{\omega}_z(0, 0, 1)^T$ which is constant in time. Moreover, all quantities are invariant along the vertical direction, so we can directly consider a

scalar velocity $\hat{\omega} : \hat{\Omega} \rightarrow \mathbb{R}$ ($\hat{\omega}$ is the velocity of the fluid on $\hat{\Omega}_f$ and of the solid in $\hat{\Omega}_s$), see Figure 2.

The constitutive equations write

$$\begin{cases} \operatorname{div} \hat{\tau} = p_z + \rho_f g & \text{on } \hat{\Omega}_f, \\ \operatorname{div} \hat{\tau} = p_z \text{ and } \nabla \hat{\omega} = 0 & \text{on } \hat{\Omega}_s, \end{cases} \quad (1)$$

with p_z the pressure gradient along the vertical direction, and where $\hat{\tau}$ is the stress, which on $\hat{\Omega}_f$ is related to the velocity by the von Mises criterion

$$\begin{cases} \hat{\tau} = \left(\mu_f + \frac{\tau_Y}{|\nabla \hat{\omega}|} \right) \nabla \hat{\omega} & \text{if } \hat{\tau} \geq \tau_Y, \\ \nabla \hat{\omega} = 0 & \text{if } \hat{\tau} \leq \tau_Y. \end{cases} \quad (2)$$

These equations state that as long as a certain stress is not reached, there is no response of the fluid (see Figure 1).

We are considering the *exchange flow* problem, meaning that we require that the total flux across the horizontal slice is zero, that is

$$\int_{\hat{\Omega}} \hat{\omega} = 0. \quad (3)$$

Mathematically, p_z can be seen as a Lagrange multiplier for (3). Moreover, we consider the following boundary conditions:

1. We assume that on the boundaries of $\hat{\Omega}$, we have a no-slip boundary condition

$$\hat{\omega} = 0 \quad \text{on } \partial \hat{\Omega}. \quad (4)$$

2. For a steady fall motion, the forces exerted by gravity and by the fluid on the solid should be in equilibrium [25]

$$\int_{\partial \hat{\Omega}_s} \hat{\tau} \cdot n = \rho_s g |\hat{\Omega}_s| \quad (5)$$

where n denotes the outward-pointing normal to $\partial \hat{\Omega}_s$. Using (5) and the divergence theorem in the second equation of (1), we get that the total pressure gradient on $\hat{\Omega}_s$ amounts to a buoyancy force

$$\int_{\hat{\Omega}_s} p_z = (\rho_s - \rho_f) g |\hat{\Omega}_s|.$$

2.1 Eigenvalue problems

Following [9, 23], we study minimizers of the functional

$$\hat{G}^\diamond(\hat{\omega}) := \begin{cases} \frac{\mu_f}{2} \int_{\hat{\Omega}_f} |\nabla \hat{\omega}|^2 + \tau_Y \int_{\hat{\Omega}_f} |\nabla \hat{\omega}| - (\rho_s - \rho_f) g \int_{\hat{\Omega}_s} \hat{\omega} & \text{if } \hat{\omega} \in \hat{H}_\diamond \\ + \infty & \text{else,} \end{cases}$$

over

$$\hat{H}_\diamond = \left\{ v \in H_0^1(\hat{\Omega}) \mid \int_{\hat{\Omega}} v = 0, \nabla v = 0 \text{ in } \hat{\Omega}_s \right\}.$$




Figure 2: Anti-plane situation: A falling cylinder, with gravity along its axis of symmetry.

Writing the Euler-Lagrange equations for \hat{G}^\diamond we obtain (1) and (2). Notice that since we work in \hat{H}_\diamond , the no-slip boundary condition (4) is automatically satisfied, and adequate testing directions are constant on connected components of $\hat{\Omega}_s$, which leads to the force balance condition (5).

We proceed to simplify the dimensions in the above functional, so that we can work with just one parameter. Assuming a given length scale \hat{L} , we define the buoyancy number Y and a velocity scale $\hat{\omega}_0$ by

$$Y := \frac{\tau_Y}{(\rho_s - \rho_f)g\hat{L}}, \quad \hat{\omega}_0 := \frac{(\rho_s - \rho_f)g\hat{L}^2}{\mu_f},$$

so that defining the rescaled velocity ω and corresponding domains by

$$\omega(x) := \frac{\hat{\omega}(\hat{L}x)}{\hat{\omega}_0}, \quad \Omega := \frac{\hat{\Omega}}{\hat{L}}, \quad \Omega_f := \frac{\hat{\Omega}_f}{\hat{L}}, \quad \text{and} \quad \Omega_s := \frac{\hat{\Omega}_s}{\hat{L}}, \quad (6)$$

we end up with the functional

$$G_Y^\diamond(\omega) := \begin{cases} \frac{1}{2} \int_{\Omega_f} |\nabla \omega|^2 + Y \int_{\Omega_f} |\nabla \omega| - \int_{\Omega_s} \omega & \text{if } \omega \in H_\diamond \\ +\infty & \text{else,} \end{cases} \quad (7)$$

over

$$H_\diamond = \left\{ v \in H_0^1(\Omega) \mid \int_{\Omega} v = 0, \quad \nabla v = 0 \text{ in } \Omega_s \right\}.$$

By the direct method it is easy to prove (see for instance [9]) that G_Y^\diamond has a unique minimizer, which corresponds to the weak solution in physical dimensions through the transformations in (6). Denoting this minimizer by ω_Y we have that for every $v \in H_\diamond$,

$$\int_{\Omega_f} \nabla \omega_Y \cdot \nabla (v - \omega_Y) + Y \int_{\Omega_f} |\nabla v| - Y \int_{\Omega_f} |\nabla \omega_Y| \geq \int_{\Omega_s} (v - \omega_Y). \quad (8)$$

As in [9], one can introduce

$$Y_c := \sup_{\omega \in H_\diamond} \frac{\int_{\Omega_s} \omega}{\int_{\Omega} |\nabla \omega|} \quad (9)$$

and test inequality (8) with $v = 0$ and $v = 2\omega_Y$ to obtain

$$\int_{\Omega} |\nabla \omega_Y|^2 = \int_{\Omega_f} |\nabla \omega_Y|^2 = \int_{\Omega_s} \omega_Y - Y \int_{\Omega_f} |\nabla \omega_Y|. \quad (10)$$

From this, and using the definition of Y_c in (9) it follows that

$$\int_{\Omega} |\nabla \omega_Y|^2 \leq \int_{\Omega_f} |\nabla \omega_Y| \left[\sup_{\omega \in H_{\diamond}} \frac{\int_{\Omega_s} \omega}{\int_{\Omega} |\nabla \omega|} - Y \right] = (Y_c - Y) \int_{\Omega_f} |\nabla \omega_Y|.$$

The last inequality implies, thanks to the homogeneous boundary conditions on ω , that $\omega_Y = 0$ in Ω_f as soon as $Y \geq Y_c$.

3 Relaxed problem and physical meaning

We determine the critical yield stress Y_c , defined in (9) and properties of the associated eigenfunction. The optimization problem (9) is equivalent to computing minimizers of the functional

$$E(\omega) := \frac{\int_{\Omega} |\nabla \omega|}{\int_{\Omega_s} \omega} \text{ over } H_{\diamond}. \quad (11)$$

Because E might not attain a minimizer in H_{\diamond} , we consider a relaxed formulation on a subset of functions of bounded variation.

3.1 Functions of bounded variations and their properties

We recall the definition of the space of functions of bounded variation and some properties of such functions that we will use below. Proofs and further results can be found in [2], for example.

Definition 1. *Let $A \subset \mathbb{R}^2$ be open. A function $v \in L^1(A)$ is said to be of bounded variation if its distributional gradient ∇v is a Radon measure with finite mass, which we denote by $\text{TV}(v)$. In particular, if $\nabla v \in L^1(A)$, then $\text{TV}(v) = \int_A |\nabla v|$. Similarly, for a set B with finite Lebesgue measure $|B| < +\infty$ we define its perimeter to be the total variation of its characteristic function 1_B , that is, $\text{Per}(B) = \text{TV}(1_B)$.*

Theorem 1. *The space of functions of bounded variation on A , denoted $\text{BV}(A)$, is a Banach space when associated with the norm*

$$\|v\|_{\text{BV}(A)} := \|v\|_{L^1(A)} + \text{TV}(v).$$

The space of functions of bounded variation satisfies the following compactness property [2, Theorem 3.44]:

Theorem 2 (Compactness and lower semi-continuity in BV). *Let $v_n \in \text{BV}(A)$ be a sequence of functions such that $\|v_n\|_{\text{BV}(A)}$ is bounded. Then there exists $v \in \text{BV}(A)$ for which, possibly upon taking a subsequence, we have*

$$v_n \xrightarrow{L^1} v.$$

In addition, for any sequence (w_n) that converges to some w in L^1 ,

$$\text{TV}(w) \leq \liminf \text{TV}(w_n).$$

We frequently use the *coarea* and *layer cake formulas*:

Lemma 1. *Let $u \in \text{BV}(\mathbb{R}^2)$ with compact support, then the coarea formula [2, Theorem 3.40]*

$$\text{TV}(u) = \int_{-\infty}^{\infty} \text{Per}(u > t) dt = \int_{-\infty}^{\infty} \text{Per}(u < t) dt \quad (12)$$

holds. If $u \in L^1(\mathbb{R}^2)$ is non-negative, then we also have the layer cake formula [19, Theorem 1.13]

$$\int_{\mathbb{R}^2} u = \int_0^{\infty} |\{u > t\}| dt. \quad (13)$$

An important role in characterizing constrained minimizers of the TV functional is played by Cheeger sets, which we now define.

Definition 2. *A set is called Cheeger set of $A \subseteq \mathbb{R}^2$ if it minimizes the ratio $\text{Per}(\cdot)/|\cdot|$ among the subsets of A .*

The following result is well known and has been stated for instance in [18, Proposition 3.5, iii] and [22, Proposition 3.1]:

Theorem 3. *For every non-empty measurable set $A \subseteq \mathbb{R}^2$ open, there exists at least one Cheeger set, and its characteristic function minimizes the quotient $u \mapsto \text{TV}(u)/\|u\|_{L^1(A)}$ in $L^1(A) \setminus \{0\}$. Moreover, almost every level set of every minimizer of this quotient is a Cheeger set.*

Remark 1. Some sets may have more than one Cheeger set, which introduces nonuniqueness in the minimizers of the quotient $\text{TV}(\cdot)/\|\cdot\|_{L^1(A)}$. One example is the set Ω of Figure 6 below.

3.2 Generalized minimizers of E

Using the compactness Theorem 2, it follows that the relaxed quotient

$$E(\omega) := \frac{\text{TV}(\omega)}{\int_{\Omega_s} \omega}$$

of (11) attains a minimizer in the space

$$\mathcal{B} := \left\{ v \in \text{BV}(\mathbb{R}^2) \mid \int_{\Omega} v = 0, \nabla v = 0 \text{ on } \Omega_s, v = 0 \text{ on } \mathbb{R}^2 \setminus \Omega \right\}.$$

Note that the quotient E is invariant with respect to scalar multiplication, and we can therefore add the constraint

$$\int_{\Omega_s} v := \frac{1}{|\Omega_s|} \int_{\Omega_s} v = 1 \quad (14)$$

to \mathcal{B} without changing the minimal value of the functional E . Thus, the problem of minimizing E over \mathcal{B} is equivalent to the following problem:

Problem 1. *Find a minimizer of TV over the set*

$$\text{BV}_{\diamond} := \left\{ v \in \text{BV}(\mathbb{R}^2) \mid \int_{\Omega} v = 0, \int_{\Omega_s} v = 1, \nabla v = 0 \text{ on } \Omega_s, v = 0 \text{ on } \mathbb{R}^2 \setminus \Omega \right\}.$$

By using standard compactness and lower semicontinuity results in $\text{BV}(\mathbb{R}^2)$, it is easy to see [9] that there is at least one solution to Problem 1. In particular, we emphasize that all the constraints above are closed with respect to the L^1 topology.

Remark 2. Notice that BV_{\diamond} is larger than the optimization space (27) used in [9], where it has been assumed that $v = \text{const.}$ in Ω_s . See also Section 5.3.

3.3 The critical yield limit

We investigate the limit of ω_Y (the minimizer of \mathcal{G}_Y^\diamond , defined in (7)) when $Y \rightarrow Y_c$. For this purpose we first prove

Proposition 1. *The quantity $\int_{\Omega_f} |\nabla \omega_Y|$ is nonincreasing with respect to $0 \leq Y \leq Y_c$. In particular, it is bounded.*

Proof. Let $Y_c \geq Y_1 > Y_2 \geq 0$. Then, from the definition (7) of ω_Y being a minimizer of \mathcal{G}_Y^\diamond it follows that

$$\mathcal{G}_{Y_2}^\diamond(\omega_{Y_2}) \leq \mathcal{G}_{Y_2}^\diamond(\omega_{Y_1}) = \mathcal{G}_{Y_1}^\diamond(\omega_{Y_1}) + (Y_2 - Y_1) \int_{\Omega_f} |\nabla \omega_{Y_1}|,$$

$$\mathcal{G}_{Y_1}^\diamond(\omega_{Y_1}) \leq \mathcal{G}_{Y_1}^\diamond(\omega_{Y_2}) = \mathcal{G}_{Y_2}^\diamond(\omega_{Y_2}) + (Y_1 - Y_2) \int_{\Omega_f} |\nabla \omega_{Y_2}|,$$

and summing, we get

$$(Y_1 - Y_2) \left(\int_{\Omega_f} |\nabla \omega_{Y_2}| - \int_{\Omega_f} |\nabla \omega_{Y_1}| \right) \geq 0,$$

which implies the assertion. \square

We are now ready to investigate the convergence of ω_Y and its rate.

Theorem 4. *For $Y \nearrow Y_c$ and any $\epsilon > 0$, we have*

$$\int_{\Omega} |\nabla \omega_Y|^2 = o((Y_c - Y)^{2-\epsilon}). \quad (15)$$

Moreover, the sequence of rescaled profiles

$$v_Y := \frac{\omega_Y}{\int_{\Omega} |\nabla \omega_Y|} \quad (16)$$

converges in the sense of Theorem 2, up to possibly taking a sequence, to a solution of Problem 1.

Proof. Using (10), the definition of Y_c (9) and since $\nabla \omega_Y$ vanishes on Ω_s , and $\int_{\Omega} |\nabla \omega_Y|$ is bounded by Proposition 1 we get

$$0 \leq \int_{\Omega_s} \omega_Y - Y \int_{\Omega} |\nabla \omega_Y| = \int_{\Omega} |\nabla \omega_Y|^2 \leq (Y_c - Y) \int_{\Omega} |\nabla \omega_Y| \leq C(Y_c - Y), \quad (17)$$

where here and in the following we denote by C a generic constant, which may be different in each appearance.

That is, we get in particular

$$\int_{\Omega} |\nabla \omega_Y|^2 \leq C(Y_c - Y). \quad (18)$$

The rate in this estimate can be improved. To do so, we use Poincaré's inequality, which combined with (18) implies that

$$\int_{\Omega_s} \omega_Y \leq C \|\omega_Y\|_{L^2(\Omega)} \leq C \sqrt{\int_{\Omega} |\nabla \omega_Y|^2} \leq C(\sqrt{Y_c - Y}). \quad (19)$$

Now, using (19) and (18) in (17) we get

$$0 \leq C(\sqrt{Y_c - Y}) - Y \int_{\Omega} |\nabla \omega_Y| \leq C(Y_c - Y),$$

which can only hold as $Y \rightarrow Y_c$ if

$$\int_{\Omega} |\nabla \omega_Y| = \mathcal{O}(\sqrt{Y_c - Y}). \quad (20)$$

But then, using (20) in (17) provides us with $\int_{\Omega} |\nabla \omega_Y|^2 = \mathcal{O}((Y_c - Y)^{3/2})$. Iterating the above substitutions using the improved estimates for $\int_{\Omega} |\nabla \omega_Y|^2$ instead of (18) (that is, using a bootstrapping argument) we obtain (15) for any $\epsilon > 0$.

Now, the associated functions v_Y , defined in (16), have total variation 1 and zero mean. From Theorem 2 it follows that v_Y converges in L^1 to some v_c . Now, it follows directly from (17) that

$$\lim_{Y \rightarrow Y_c} \frac{Y \int_{\Omega} |\nabla \omega_Y|}{\int_{\Omega_s} \omega_Y} = 1, \quad (21)$$

and therefore, using the L^1 convergence of v_Y , its definition (16) and that $\int_{\Omega} |\nabla \omega_y| = \int_{\Omega_f} |\nabla \omega_y|$, (21) implies

$$\int_{\Omega_s} v_c = \lim_{Y \rightarrow Y_c} \int_{\Omega_s} v_Y = \lim_{Y \rightarrow Y_c} \frac{\int_{\Omega_s} \omega_Y}{\int_{\Omega_f} |\nabla \omega_y|} = Y_c.$$

Recalling that $\text{TV}(v_Y) = 1$, the semi-continuity of the total variation with respect to L^1 convergence implies $\text{TV}(v_c) \leq 1$, which yields

$$Y_c \int_{\Omega} |\nabla v_c| - \int_{\Omega_s} v_c \leq 0,$$

which can be rewritten as

$$Y_c \leq \frac{\int_{\Omega_s} v_c}{\int_{\Omega} |\nabla v_c|}$$

so v_c is a maximizer of $v \mapsto \frac{\int_{\Omega_s} v}{\int_{\Omega} |\nabla v|}$. □

From the above result, we see that a minimizer of the quotient $\frac{\int_{\Omega} |\nabla v|}{\int_{\Omega_s} v}$ can be obtained as a limit of rescaled physical velocities, and therefore carries information about their geometry. For this reason, we will focus on these minimizers in the following.

4 Piecewise constant minimizers

We prove the existence of solutions of Problem 1 with particular properties. In our previous work [9] this problem was considered under the assumption that the velocity is constant in the whole Ω_s . In the situation considered here, the physical velocity ω is constant only on every connected component of Ω_s , and the velocity of *each* solid particle is an unknown. Therefore, the candidates of limiting profiles v over which we optimize (belonging to BV_{\diamond}) also satisfy $\nabla v = 0$ on Ω_s .

4.1 A minimizer with three values

Theorem 5. *There is a solution of Problem 1 that attains only two non-zero values.*

The same result has been proved in [9] in the simpler situation when the velocities were considered uniformly constant on the whole Ω_s . For the proof of Theorem 5, we proceed in two steps:

1. We prove the existence of a minimizer for Problem 1 with attains only finitely many values. This is accomplished by convexity arguments reminiscent of slicing by the coarea (12) and layer cake (13) formulas, but more involved.
2. When considered over functions with finitely many values, the minimization of the total variation with integral constraints is a simple finite-dimensional optimization problem, and standard linear programming arguments provide the result.

Step 1. A minimizer with finite range. To begin the proof, we assume that we are given a minimizer u of the total variation in BV_\diamond , that is, a solution of Problem 1. We represent Ω_s by its connected components Ω_s^i , $i = 1, \dots, N$,

$$\Omega_s = \bigcup_{i=1}^N \Omega_s^i.$$

Since u belongs to BV_\diamond , u is constant on every Ω_s^i , and we introduce the constants γ_i such that

$$u|_{\Omega_s^i} = \gamma_i.$$

Note that the constraint (14) reads

$$\frac{1}{\sum_{i=1}^n |\Omega_s^i|} \sum_{i=1}^N \gamma_i |\Omega_s^i| = 1.$$

Defining

$$u_i := u \cdot 1_{\{\gamma_i < u < \gamma_{i+1}\}} + \gamma_i 1_{\{u \leq \gamma_i\}} + \gamma_{i+1} 1_{\{u \geq \gamma_{i+1}\}},$$

we have

$$u = \sum_{i=1}^N (u_i - \gamma_i).$$

Notice that each u_i minimizes the total variation among functions with fixed integral $\int_\Omega u_i$, and satisfying the boundary conditions $u = \gamma_i$ on $\{u \leq \gamma_i\}$ and $u = \gamma_{i+1}$ on $\{u \geq \gamma_{i+1}\}$.

As a result, the function $v_i := \frac{u_i - \gamma_i}{\gamma_{i+1} - \gamma_i}$ minimizes the total variation with constraints $v_i|_{\mathbb{R}^2 \setminus \{u > \gamma_i\}} \equiv 0$, $v_i|_{\{u \geq \gamma_{i+1}\}} \equiv 1$ and prescribed integral. Lemma 2 below (applied with $\Omega_0 = \{u > \gamma_i\}$ and $\Omega_1 = \{u \geq \gamma_{i+1}\}$) shows that v_i can be replaced by a five level-set function \tilde{v}_i which has total variation smaller or equal to $\text{TV}(v_i)$. Hence u_i can be replaced by the five level-set function $\tilde{u}_i := \gamma_i + \tilde{v}_i(\gamma_{i+1} - \gamma_i)$ without increasing the total variation.

Therefore, the finitely-valued function

$$\tilde{u} := \sum_{i=1}^N (\tilde{u}_i - \gamma_i)$$

is again a solution of Problem 1 (the functions u and \tilde{u} coincide on Ω_s , so the constraint $\int_{\Omega_s} \tilde{u} = 1$ is satisfied).

We complete Step 1 by proving Lemma 2, which again requires the proof of an auxiliary lemma.

Lemma 2. *Let $\Omega_1 \subset \Omega_0$ be two bounded measurable sets, $\nu \in \mathbb{R}$. Then, there exists a minimizer of TV on the set*

$$\mathcal{A} := \left\{ v \in \text{BV}(\mathbb{R}^2) \mid v|_{\mathbb{R}^2 \setminus \Omega_0} \equiv 0, v|_{\Omega_1} \equiv 1, \int_{\mathbb{R}^2} v = \nu \right\},$$

where the range consists of at most five values, one of them being zero.

Proof. Let w be an arbitrary minimizer of TV in \mathcal{A} . Splitting w at 0 and 1 we can write

$$w = (w^{1+} - 1) + w^{(0,1)} - w^- \quad (22)$$

with $w^{1+} := w \cdot 1_{w \geq 1} + 1_{w < 1}$, $w^{(0,1)} = w \cdot 1_{0 \leq w < 1} + 1_{w > 1}$, and w^- the usual negative part. We see from the coarea formula that

$$\begin{aligned} \text{TV}(w) &= \int_{s \leq 0} \text{Per}(w \geq s) + \int_{0 < s < 1} \text{Per}(w \geq s) + \int_{s \geq 1} \text{Per}(w \geq s) \\ &= \text{TV}(w^-) + \text{TV}(w^{(0,1)}) + \text{TV}(w^{1+}). \end{aligned}$$

With this splitting, w^- can be seen to be a minimizer of TV over

$$\mathcal{A}^- := \left\{ v \in \text{BV}(\mathbb{R}^2) \mid v = 0 \text{ on } \{w > 0\} \cup \mathbb{R}^2 \setminus \Omega_0, \int_{\Omega} v = \int_{\Omega} w^- \right\}.$$

By Theorem 3, almost every level set of w^- is a Cheeger set of $\Omega_0 \setminus \{w > 0\}$, the complement of $\{w > 0\} \cup \mathbb{R}^2 \setminus \Omega_0$. In particular, if we replace w^- by $\frac{\int_{\Omega} w^-}{|\mathcal{C}_0|} 1_{\mathcal{C}_0}$, where \mathcal{C}_0 is one such Cheeger set, the total variation doesn't increase. Therefore, there exists a minimizer \tilde{w}^- of TV on \mathcal{A}^- that reaches only one non-zero value.

With an analogous argumentation we see that, because w^{1+} minimizes TV on the set

$$\mathcal{A}^{1+} := \left\{ v \in \text{BV}(\mathbb{R}^2) \mid v = 1 \text{ on } \{w < 1\}, \int_{\Omega} v = \int_{\Omega} w^{1+} \right\},$$

there exists a minimizer \tilde{w}^{1+} that writes

$$\tilde{w}^{1+} = 1 + \zeta 1_{\mathcal{C}_1}$$

where \mathcal{C}_1 is a Cheeger set of $\{w \geq 1\}$ and $\zeta \geq 0$ is a constant.

Moreover, defining

$$\mu := \int_{\Omega} w^{(0,1)},$$

$w^{(0,1)}$ minimizes TV on the set

$$\mathcal{A}_{\mu}^{(0,1)} := \left\{ v \in \text{BV}(\mathbb{R}^2) \mid v = 1 \text{ on } \{w \geq 1\}, v = 0 \text{ on } \{w \leq 0\} \text{ and } \int_{\Omega} v = \mu \right\}.$$

We now show that there exists a minimizer of TV in $\mathcal{A}_{\mu}^{(0,1)}$ that attains only three values. Since $w^{(0,1)}$ is one of them, there exists some minimizer of TV in $\mathcal{A}_{\mu}^{(0,1)}$ with values in $[0, 1]$. We denote by u a generic one. In what follows, we denote by $E_s := \{u > s\}$ the level-sets of u .

We prove in Lemma 3 that for almost every s , 1_{E_s} minimizes TV in $\mathcal{A}_{|E_s|}^{(0,1)}$. That implies in particular that for a.e. s , E_s minimizes perimeter with fixed mass. We introduce $E_s^{(1)}$ the set of points of density 1 for E_s and $E_s^{(0)}$ the set of points of density 0 for E_s , that is

$$E_s^{(1)} := \left\{ x \in \Omega \left| \lim_{r \rightarrow 0} \frac{|E_s \cap B_r(x)|}{|B_r(x)|} = 1 \right. \right\} \quad \text{and} \quad E_s^{(0)} := \left\{ x \in \Omega \left| \lim_{r \rightarrow 0} \frac{|E_s \cap B_r(x)|}{|B_r(x)|} = 0 \right. \right\}.$$

Lebesgue differentiation theorem implies that $E_s^{(1)} = E_s$ and $E_s^{(0)} = \Omega \setminus E_s$ a.e.

Now, since the level-sets are nested, the function $s \mapsto |E_s|$ is nonincreasing. Therefore, there exists s_μ such that

$$\text{for } s > s_\mu, |E_s| \leq \mu, \text{ and for } s < s_\mu, |E_s| \geq \mu.$$

Let us now define

$$E^+ := \bigcup_{s > s_\mu} E_s^{(1)} \quad \text{and} \quad E^- := \bigcap_{s < s_\mu} \Omega \setminus E_s^{(0)}.$$

We then have

Claim. *If E^\pm is not empty, 1_{E^\pm} minimizes total variation in $\mathcal{A}_{|E^\pm|}^{(0,1)}$, with $|E^+| \leq \mu \leq |E^-|$.*

Proof of claim. By Lemma 3, $1_{E_s^{(1)}}$ minimizes total variation in $\mathcal{A}_{|E_s^{(1)|}}^{(0,1)}$ for almost every s . Then, let us select a decreasing sequence $s_n \searrow s_\mu$ such that for each n , $1_{E_{s_n}^{(1)}}$ minimizes total variation in $\mathcal{A}_{|E_{s_n}^{(1)}|}^{(0,1)}$. Since $E_{s_n}^{(1)} \rightarrow E^+$ in L^1 , one has $|E^+| = \lim |E_{s_n}^{(1)}| = \lim |E_{s_n}|$ and the semicontinuity for the perimeter gives

$$\text{Per}(E^+) \leq \liminf \text{Per}(E_{s_n}^{(1)}).$$

In fact, the sequence $\text{Per}(E_{s_n}^{(1)})$ is bounded. To see this, we fix a value $\hat{s} < s_\mu$ and since $E_{s_1} \subset E_{s_n}^{(1)} \subset E_{\hat{s}}$ we can write for some $t_n \in (0, 1)$

$$|E_{s_n}^{(1)}| = t_n |E_{\hat{s}}| + (1 - t_n) |E_{s_1}|.$$

Therefore, applying Lemma 3 again we obtain

$$\text{Per}(E_{s_n}^{(1)}) \leq \text{TV}(t_n 1_{E_{\hat{s}}} + (1 - t_n) 1_{E_{s_1}}) \leq \text{Per}(E_{\hat{s}}) + \text{Per}(E_{s_1}).$$

Now, let us assume that there exists $v \in \text{BV}(\Omega)$ with $\int v = |E^+|$ and $\text{TV}(v) < \text{Per}(E^+) - \varepsilon$. By the above, for every $\delta > 0$ we can find n such that $|E^+| \geq |E_{s_n}| \geq |E^+| - \delta$ and

$$\text{Per}(E^+) \leq \text{Per}(E_{s_n}^{(1)}) + \delta.$$

Now, if $\delta < \varepsilon/10$ is small enough, we can find a ball $B_n \subset \Omega$ such that $\int_\Omega v \cdot 1_{\Omega \setminus B_n} = |E_{s_n}|$ and $\|v\|_\infty \text{Per}(B_n) \leq \varepsilon/10$, so we get

$$\begin{aligned} \text{TV}(v \cdot 1_{\Omega \setminus B_n}) &\leq \text{TV}(v) + \|v\|_\infty \text{Per}(B_n) \leq \text{Per}(E^+) - \varepsilon + \|v\|_\infty \text{Per}(B_n) \\ &\leq \text{Per}(E_{s_n}^{(1)}) + \|v\|_\infty \text{Per}(B_n) + \delta - \varepsilon \leq \text{Per}(E_{s_n}^{(1)}) - \frac{\varepsilon}{2}, \end{aligned}$$

and therefore we get a contradiction with the TV-minimality of $E_{s_n}^{(1)}$.

Selecting an increasing sequence $\tilde{s}_n \nearrow s_\mu$ and such that $\Omega \setminus E_{\tilde{s}_n}^{(0)}$ minimizes TV in $\mathcal{A}_{|E_{\tilde{s}_n}^{(0)}|}^{(0,1)}$, we obtain similarly that 1_{E^-} minimizes TV in $\mathcal{A}_{|E^-|}^{(0,1)}$. \square

To finish the proof of Lemma 2, we distinguish two alternatives. Either E^+ or E^- has mass μ , in which case the claim above implies Lemma 2, or E^\pm are both nonempty and

$$|E^+| < \mu \quad \text{and} \quad |E^-| > \mu.$$

In the second case, let $s < s_\mu$. Then, $|E^-| \in (|E^+|, |E_s|)$ and there exists $t = \frac{|E^-| - |E^+|}{|E_s| - |E^+|}$ such that $|E^-| = t|E_s| + (1-t)|E^+|$. The function $t1_{E_s} + (1-t)1_{E^+}$ therefore belongs to $\mathcal{A}_{|E^-|}^{(0,1)}$. Since 1_{E^-} is a minimizer of TV in this set, one must have

$$\text{Per}(E^-) \leq \text{TV}(t1_{E_s} + (1-t)1_{E^+}) \leq \frac{|E^-| - |E^+|}{|E_s| - |E^+|} \text{Per}(E_s) + \frac{|E_s| - |E^-|}{|E_s| - |E^+|} \text{Per}(E^+).$$

This equation rewrites

$$\text{Per}(E_s) \geq \frac{|E_s| - |E^+|}{|E^-| - |E^+|} \text{Per}(E^-) + \frac{|E^-| - |E_s|}{|E^-| - |E^+|} \text{Per}(E^+). \quad (23)$$

Similarly, if $s > s_\mu$, one has $|E_s| < |E^+|$ and $|E^+|$ is a convex combination of $\{|E^-|, |E_s|\}$. The same steps lead to the same (23). Finally, one just write (we use (23), the coarea and the layer-cake formulas)

$$\begin{aligned} \text{TV}(u) &= \int_0^1 \text{Per}(E_s) \geq \int_0^1 \frac{(|E_s| - |E^+|) \text{Per}(E^-) + (|E^-| - |E_s|) \text{Per}(E^+)}{|E^-| - |E^+|} \\ &\geq \int_0^1 \frac{\text{Per}(E^-) - \text{Per}(E^+)}{|E^-| - |E^+|} |E_s| + \frac{|E^-| \text{Per}(E^+) - |E^+| \text{Per}(E^-)}{|E^-| - |E^+|} \\ &= \frac{\text{Per}(E^-) - \text{Per}(E^+)}{|E^-| - |E^+|} \mu + \frac{|E^-| \text{Per}(E^+) - |E^+| \text{Per}(E^-)}{|E^-| - |E^+|} \\ &= \text{TV}(\lambda 1_{E^-} + (1-\lambda)1_{E^+}) \end{aligned}$$

with $\lambda = \frac{\mu - |E^+|}{|E^-| - |E^+|}$.

As a result, one can replace $w^{(0,1)}$ in the decomposition (22) by a three valued minimizer $\tilde{w}^{(0,1)}$ of TV in $\mathcal{A}_\mu^{(0,1)}$. Therefore, combining the three modified parts we see that there exists a minimizer in \mathcal{A}

$$\tilde{w} := (\tilde{w}^{1+} - 1) + \tilde{w}^{(0,1)} - \tilde{w}^-$$

which attains at most five values. □

Lemma 3. *Let u be a minimizer of TV in $\mathcal{A}_\mu^{(0,1)}$ with values in $[0, 1]$, and $E_s := \{u > s\}$. Let s_0 be a Lebesgue point of $s \mapsto \text{Per}(E_s)$ and $s \mapsto |E_s|$ (these two functions are measurable, so almost every $s \in [0, 1]$ is a Lebesgue point for them). Then $1_{E_{s_0}}$ minimizes TV in $\mathcal{A}_{|E_{s_0}|}^{(0,1)}$.*

Proof. First, note that for every $s_1 < s_2$, the function

$$u_{[s_1, s_2]} := s_2 1_{E_{s_2}} + u \cdot 1_{[s_1, s_2]} + s_1 1_{u < s_1}$$

is such that $v := \frac{u_{[s_1, s_2]} - s_1}{s_2 - s_1}$ minimizes the total variation in $\mathcal{A}_{\int v}^{(0,1)}$. Indeed, if $\hat{v} \in \mathcal{A}_{\int v}^{(0,1)}$ with $\text{TV}(\hat{v}) < \text{TV}(v)$, then $\text{TV}(\hat{v}(s_2 - s_1) + s_1) < \text{TV}(u_{[s_1, s_2]})$. Since $u = (u \cdot 1_{u < s_1} - s_1) + u_{[s_1, s_2]} + (u \cdot 1_{u > s_2} - s_2)$, then we would have

$$\begin{aligned} \text{TV}(u) &= \text{TV}((u \cdot 1_{u < s_1} - s_1)) + \text{TV}(u_{[s_1, s_2]}) + \text{TV}((u \cdot 1_{u > s_2} - s_2)) \\ &> \text{TV}((u \cdot 1_{u < s_1} - s_1)) + \text{TV}(\hat{v}(s_2 - s_1) + s_1) + \text{TV}((u \cdot 1_{u > s_2} - s_2)) \\ &\geq \text{TV}((u \cdot 1_{u < s_1} - s_1) + (\hat{v}(s_2 - s_1) + s_1) + (u \cdot 1_{u > s_2} - s_2)), \end{aligned}$$

where $(u \cdot 1_{u < s_1} - s_1) + (\hat{v}(s_2 - s_1) + s_1) + (u \cdot 1_{u > s_2} - s_2) \in \mathcal{A}_\mu^{(0,1)}$, which is a contradiction with the minimality of u .

Letting s_0 as in the assumptions, we have just seen that for every $h > 0$, $\frac{u_{[s_0-h, s_0+h]} - (s_0-h)}{2h}$ minimizes the total variation in $\mathcal{A}_{\mu_h}^{(0,1)}$ with

$$\begin{aligned} \mu_h &:= \frac{\int_{s_0-h \leq u \leq s_0+h} (u - (s_0 - h))}{2h} + |E_{s_0+h}| = \frac{1}{2h} \int_{s_0-h}^{s_0+h} |\{u > t\} \cap \{u \leq s_0 + h\}| dt + |E_{s_0+h}| \\ &= \frac{1}{2h} \int_{s_0-h}^{s_0+h} |\{u > t\}| dt = \frac{1}{2h} \int_{s_0-h}^{s_0+h} |E_s| ds. \end{aligned}$$

On the other hand, the total variation of $\frac{u_{[s_0-h, s_0+h]} - (s_0-h)}{2h}$ writes, using the coarea formula,

$$\frac{1}{2h} \int_{s_0-h}^{s_0+h} \text{Per}(E_s).$$

Finally, let us assume that $1_{E_{s_0}}$ does not minimize total variation in $\mathcal{A}_{|E_{s_0}|}^{(0,1)}$. Then, there would exist $\varepsilon > 0$ and $u_0 \in \mathcal{A}_{|E_{s_0}|}^{(0,1)}$ such that

$$\text{Per}(E_{s_0}) \geq \text{TV}(u_0) + \varepsilon.$$

Since s_0 is a Lebesgue point, one can find $\delta > 0$ such that for every $h \leq \delta$,

$$\left| \frac{1}{2h} \int_{s_0-h}^{s_0+h} \text{Per}(E_s) ds - \text{Per}(E_{s_0}) \right| \leq \frac{\varepsilon}{10} \quad \text{and} \quad \left| \frac{1}{2h} \int_{s_0-h}^{s_0+h} |E_s| ds - |E_{s_0}| \right| \leq \frac{\varepsilon}{10}.$$

Let $h \leq \delta$ and B be a ball such that $\text{Per}(B) \leq \frac{\varepsilon}{4\|u_0\|_\infty}$. There exists α such that the function $u_0 + \alpha 1_B$ satisfies

$$\int u_0 + \alpha 1_B = \frac{1}{2h} \int_{s_0-h}^{s_0+h} |E_s| ds.$$

Reducing h if needed, one can enforce that $|\alpha| \leq 2\|u_0\|_\infty$.

Then,

$$\text{TV}(u_0 + \alpha 1_B) \leq \text{TV}(u_0) + \alpha \text{Per}(B) \leq \text{Per}(E_{s_0}) - \varepsilon + \alpha \text{Per}(B) \leq \frac{1}{2h} \int_{s_0-h}^{s_0+h} \text{Per}(E_s) ds - \frac{4\varepsilon}{10},$$

which contradicts the minimality of $u_{[s_0-h, s_0+h]}$ and proves the claim. \square

Step 2. Construction of a three-valued minimizer. Step 1 provides a solution \tilde{u} of Problem 1 that reaches $p+1$ values. We denote its range by

$$\{\gamma_{p^-}, \dots, \gamma_{-1}, 0, \gamma_1, \dots, \gamma_{p^+}\}$$

where $p^- \leq 0 \leq p^+$, $p^+ - p^- = p$ and $\gamma_i < 0$ for $i < 0$ and $\gamma_i > 0$ for $i > 0$.

Let us now define, for $i < 0$, $E_i := \{\tilde{u} \leq \gamma_i\}$ and $\alpha_i := \gamma_i - \gamma_{i+1}$ and for $i > 0$, $E_i := \{\tilde{u} \geq \gamma_i\}$ and $\alpha_i := \gamma_i - \gamma_{i-1}$. The function \tilde{u} then writes

$$\tilde{u} = \sum_{\substack{i=p^- \\ i \neq 0}}^{p^+} \alpha_i 1_{E_i} \tag{24}$$

where $E_i \subset E_j$ whenever $i < j < 0$ or $i > j > 0$.

We also have

$$\begin{aligned} \text{TV}(\tilde{u}) &= \sum_{\substack{i=p^- \\ i \neq 0}}^{p^+} |\alpha_i| \text{Per}(E_i), \\ \int_{\Omega} \tilde{u} &= \sum_{\substack{i=p^- \\ i \neq 0}}^{p^+} \alpha_i |E_i|, \\ \int_{\Omega_s} \tilde{u} &= \sum_{\substack{i=p^- \\ i \neq 0}}^{p^+} \alpha_i |E_i^s| \end{aligned} \tag{25}$$

where $E_i^s = E_i \cap \Omega_s$.

Since \tilde{u} is a solution to Problem 1, the collection (α_i) minimizes $\sum_i |\alpha_i| \text{Per}(E_i)$ with constraints

$$\sum_{\substack{i=p^- \\ i \neq 0}}^{p^+} \alpha_i |E_i| = 0 \quad \text{and} \quad \sum_{\substack{i=p^- \\ i \neq 0}}^{p^+} \alpha_i |E_i^s| = |\Omega_s|$$

as well as $\alpha_i < 0$ for $i < 0$ and $\alpha_i > 0$ for $i > 0$. The constraint on the sign of the α_i is made such that the formula (25) holds. Indeed, if the α_i change signs, the right hand side of (25) is only an upper bound for $\text{TV}(\tilde{u})$.

Introducing the vectors

$$\begin{aligned} a &= (\text{Per}(E_{p^-}), \dots, \text{Per}(E_{p^+})), \\ b &= (|E_{p^-}^s|, \dots, |E_{p^+}^s|), \\ c &= (|E_{p^-}|, \dots, |E_{p^+}|), \\ x &= (\alpha_{p^-}, \dots, \alpha_{p^+}), \end{aligned}$$

minimizing (25) for \tilde{u} of the form (24) and with the constraints mentioned above is reformulated into finding a minimizer of

$$\begin{aligned} (a, x) &\rightarrow |a^T x|_{\ell^1}, \\ x \text{ s.t. } &b^T x = |\Omega_s| \text{ and } c^T x = 0. \end{aligned}$$

Denoting by $\sigma \in \{-1, 1\}^p \subseteq \mathbb{R}^p$ indexed by $i \in \{p^-, \dots, p^+\}$ with $\sigma_i = -1$ for $i < 0$ and $\sigma_i = 1$ for $i > 0$, this minimization problem can be rewritten as

$$\min_{x \in \mathbb{R}^{p+1}} \{a^T(\sigma : x) = (\sigma : a)^T x \mid b^T x = |\Omega_s|, c^T x = 0, \sigma : x \geq 0\}, \tag{26}$$

where $\sigma : x := (x_1 \sigma_1, \dots, x_p \sigma_p, x_{p+1} \sigma_{p+1})$. The space of constraints is then a (possibly empty) polyhedron given by the intersection of the quadrant $\sigma : x \geq 0$ with the two hyperplanes $c^T x = 0$ and $b^T x = |\Omega_s|$. Now for a point of a polyhedron in \mathbb{R}^p to be a vertex, we must have that at least p constraints are active at it. Therefore, at least $p - 2$ of these constraints should be of those defining the quadrant $\sigma : x \geq 0$, meaning that at a vertex, at least $p - 2$ coefficients of x are zero.

This polyhedron could be unbounded, but since $a \geq 0$ and $\sigma : x \geq 0$ componentwise, the minimization of $a^T(\sigma : x)$ must have at least one solution in it. Moreover, since it is contained in a quadrant ($\sigma : x \geq 0$), it clearly does not contain any line, so it must have at least one vertex ([5, Theorem 2.6]). Since the function to minimize is linear in x , it has a minimum at

one such vertex ([5, Theorem 2.7]). That proves the existence of a minimizer of (26) with at least $p - 2$ of the (α_i) being zero. This corresponds to a minimizer for Problem 1 which has only two level-sets with nonzero values, finishing the proof of Theorem 5.

4.2 Minimizers with connected level-sets

In this subsection, we refine our analysis slightly, and show the existence of three-valued minimizers for Problem 1 with additional properties. We start with the following definition:

Definition 3. *A set of finite perimeter A is called indecomposable, if there are no two disjoint finite perimeter sets B, C such that $|B| > 0$, $|C| > 0$, $A = B \cup C$ and $\text{Per}(A) = \text{Per}(B) + \text{Per}(C)$.*

This notion is in fact a natural measure-theoretic sense of connectedness for sets for finite perimeter, for more information about it see [1].

Remark 3. By computing the Fenchel dual of Problem 1, it can be seen that the non-zero level-sets of any solution are minimizers of the functional

$$E \mapsto \text{Per}(E) - \int_{\Omega \setminus \Omega_s} k, \text{ with } k \in L^2(\Omega \setminus \Omega_s).$$

This optimality property in turn implies lower bounds only depending on k for the perimeter and mass of E , and in case it can be decomposed in the sense of Definition 3, the same lower bounds also hold for each set in such a decomposition. In consequence, E can only be decomposed in at most a finite number of sets. The proof of these statements relies heavily on the results of [1], and is presented in [6] for the unconstrained case, and [14] for the case with Dirichlet constraints, as used here.

Assuming these results, one can simplify the level sets of solutions further:

Theorem 6. *There exists a minimizer for Problem 1 attaining exactly three values for which all non-zero level-sets are indecomposable.*

Proof. First, we consider the positive level-set and assume that it is decomposable in two sets Ω_1, Ω_2 as in Definition 3. Then the corresponding minimizer u can be written as

$$u = \alpha(1_{\Omega_1} + 1_{\Omega_2}) - \beta 1_{\Omega_-},$$

where $\alpha, \beta > 0$. Consider a perturbation of u of the form

$$u_h = (\alpha + h)1_{\Omega_1} + (\alpha + k)1_{\Omega_2} - (\beta + l)1_{\Omega_-},$$

with $|h| \leq \alpha$, $|k| \leq \alpha$, and $|l| \leq \beta$. Then, since $\Omega_1 \cap \Omega_2 = \emptyset$, $u_h \in \text{BV}_\diamond$ if and only if

$$h|\Omega_1| + k|\Omega_2| - l|\Omega_-| = 0 \quad \text{and} \quad h|\Omega_1^s| + k|\Omega_2^s| - l|\Omega_-^s| = 0,$$

where $\Omega_i^s := \Omega_i \cap \Omega_s$. These two equations lead to

$$l = h \frac{|\Omega_1^s||\Omega_2| - |\Omega_1||\Omega_2^s|}{|\Omega_2||\Omega_-^s| - |\Omega_-||\Omega_2^s|} \quad \text{and} \quad k = h \frac{|\Omega_1^s||\Omega_-| - |\Omega_1||\Omega_-^s|}{|\Omega_2||\Omega_-^s| - |\Omega_-||\Omega_2^s|}.$$

Under our assumptions on h, k, l, Ω_1 and Ω_2 , and since $1_{\Omega_1} + 1_{\Omega_2} = 1_{\Omega_1 \cup \Omega_2}$, the total variation of the perturbed function u_h can be written as

$$\begin{aligned} \text{TV}(u_h) &= (\alpha + \min(h, k)) \text{Per}(\Omega_1 \cup \Omega_2) \\ &\quad + (h - k)^+ \text{Per}(\Omega_1) + (k - h)^+ \text{Per}(\Omega_2) + (\beta + l) \text{Per}(\Omega_-) \\ &= (\alpha + h) \text{Per}(\Omega_1) + (\alpha + k) \text{Per}(\Omega_2) + (\beta + l) \text{Per}(\Omega_-). \end{aligned}$$

Then, because u is a minimizer of TV, it follows that

$$h \operatorname{Per}(\Omega_1) + k \operatorname{Per}(\Omega_2) + l \operatorname{Per}(\Omega_-) \geq 0.$$

Since the left hand side and k, l are linear in h , one can replace h by $-h$ and obtain

$$h \operatorname{Per}(\Omega_1) + k \operatorname{Per}(\Omega_2) + l \operatorname{Per}(\Omega_-) = 0$$

which shows that u_h is also a minimizer. Now since we have

$$\beta = \alpha \frac{|\Omega_1| + |\Omega_2|}{|\Omega_-|} \text{ and } l = \frac{h|\Omega_1| + k|\Omega_2|}{|\Omega_-|},$$

one can choose h such that $h = -\alpha$ or $k = -\alpha$ without violating $|l| \leq \beta$, and therefore produce a minimizer whose positive part is either Ω_2 or Ω_1 , respectively. We proceed similarly for the negative part and therefore obtain an indecomposable negative level-set. \square

Remark 4. In the above proof, through an adequate choice of components for deletion, one can even obtain simply connected level sets. For example, assuming that Ω_2 is fully enclosed in Ω_- (that is if $\partial\Omega_2 \cap \partial\Omega_- = \partial\Omega_2$), then the variation of u_h can also be written

$$\operatorname{TV}(u_h) = (\alpha + h) \operatorname{Per}(\Omega_1) + (\alpha + k + \beta + l) \operatorname{Per}(\Omega_2) + (\beta + l)(\operatorname{Per}(\Omega_-) - \operatorname{Per}(\Omega_2)),$$

which is linear in h as long as $k \geq -\alpha - \beta - l$. The equality case in this last constraint corresponds to joining Ω_2 to Ω_- , and avoiding creating a “hole” in Ω_- by the procedure mentioned above (which replaces $\alpha 1_{\Omega_2}$ by zero). Clearly, this procedure can also be performed for the positive level set, and in fact the “holes” to be deleted could also be connected components of the zero level set. Therefore, a solution in which both the positive and negative level set are simply connected can be obtained.

Remark 5. The intuition behind these last results is that, like in the proof of Theorem 5, the constraints of the problem are linear with respect to the values, and the total variation is also linear as long as the signs of the differences of values at the interfaces do not change. In particular, the points at which the topology of the level sets changes are situations in which these signs change (that is, the values of two adjacent level sets are equal).

5 Numerical scheme and results

We now turn our attention to the numerical computation of solutions to the eigenvalue for Problem 1. At first, for simplicity, we limit ourselves to the case (considered in [9]) in which the velocities are assumed constant on the whole Ω_s . That is, the problem considered is minimization of the total variation in the space

$$\operatorname{BV}_{\diamond,1} := \left\{ u \in \operatorname{BV}(\mathbb{R}^2) \mid \int_{\Omega} u = 0, u \equiv 1 \text{ in } \Omega_s, u \equiv 0 \text{ in } \mathbb{R}^2 \setminus \Omega \right\}, \quad (27)$$

where the constraint $u \equiv 1$ in Ω_s corresponds to (14) under this simplification.

This restriction corresponds to the case in which either Ω_s is connected, so that there is only one solid particle, or all the particles are constrained to move with the same velocity. In Section 5.4 we point out the required modifications for the multi-particle case and present a variety of computed examples.

To compute a minimizer of TV in $\operatorname{BV}_{\diamond,1}$, we use a standard primal dual algorithm [8]. The constraint $\int_{\Omega} v = 0$ is enforced through a scalar Lagrange multiplier q , whereas the conditions

$v = 0$ on $\partial\Omega$ and $v = 1$ on Ω_s are encoded as indicator functions. Our discretization of choice is finite differences on a rectangular grid $\{1, \dots, m\} \times \{1, \dots, n\}$, where in this whole section, for simplicity, we assume that $n = m$ and $\Omega \Subset (0, 1)^2$. This leads to a minimization problem of the form

$$\min_{v \in X} \max_{\substack{p \in X^4 \\ q \in \mathbb{R}}} \chi_{C^n}(v) + \sum_{i,j} [(\nabla v)^{ij} \cdot p^{ij} - \chi_{\{|\cdot| \leq 1\}}(p^{ij}) - qv^{ij}]. \quad (28)$$

Here, $X = \mathbb{R}^{n^2}$ denotes the space of real-valued discrete functions on the square grid $G^n = \{1, \dots, n\} \times \{1, \dots, n\}$. Since we use Dirichlet boundary conditions, the grid encloses the physical domain. The corresponding constraint set is then

$$C^n := \{v \in X \mid v = 0 \text{ on } G^n \setminus \Omega^n, v = 1 \text{ on } \Omega_s^n\}, \quad (29)$$

where Ω^n and Ω_s^n denote the parts of the grid corresponding to Ω and Ω_s respectively (note that to correctly account for perimeter at the boundary we must have $\Omega^n \subset \{2, \dots, n-1\} \times \{2, \dots, n-1\}$). The indicator function (in the convex analysis sense) of a set A is denoted by χ_A , so that $\chi_A(x) = 0$ if $x \in A$, and $+\infty$ otherwise. ∇ stands for a suitable discrete gradient, whose choice we now discuss.

5.1 Discretization

We discretize the problem using the ‘‘upwind’’ scheme of [7] which has the advantage of carrying a high degree of isotropy. The discrete velocity is denoted by v^{ij} , and we use the signed gradient $(\nabla v)^{ij}$ introduced in [7], containing separate components for forward and backward differences with opposite signs:

$$\begin{aligned} (\nabla v)^{ij} &:= (v^{i+1,j} - v^{i,j}, v^{i-1,j} - v^{i,j}, v^{i,j+1} - v^{i,j}, v^{i,j-1} - v^{i,j}) \\ &=: \left((\nabla v)_{1,+}^{ij}, (\nabla v)_{1,-}^{ij}, (\nabla v)_{2,+}^{ij}, (\nabla v)_{2,-}^{ij} \right) \end{aligned} \quad (30)$$

therefore, at each grid point $(i, j) \in G^n$ the signed gradient and its corresponding multiplier variables $\nabla v^{ij}, p^{ij} \in (\mathbb{R}^2)^2$. We note that to compute the gradient when any of the indices is 1 or n one needs to extend the functions outside the grid, but for the problem at hand any choice will do, since Ω^n never touches the boundary of the grid.

For us it is important to use a discretization that takes into account derivatives in all coordinate directions equally, since we aim to resolve sharp geometric interfaces that are not induced by a regularization data term. Figure 3 contains a comparison with the results obtained when using forward differences. In that case, the geometry of the interfaces is distorted according to their orientations, a phenomenon which is minimized in the upwind scheme. Using centered differences is also not adequate, since the centered difference operator has a nontrivial kernel and our solutions are constant in large parts of the domain.

5.2 Convergence of the discretization

It is well-known that the standard finite difference discretizations of the total variation converge, in the sense of Γ -convergence with respect to the L^1 topology [7], where the discrete functionals are appropriately defined for piecewise constant functions. We now aim to demonstrate that the chosen discretization and penalization scheme still converges and correctly accounts for the boundary conditions in the limit. We introduce, for each $(i, j) \in \{1, \dots, n-1\}^2$,

$$R_{ij}^n := \frac{1}{n} \left(i - \frac{1}{2}, i + \frac{1}{2} \right) \times \left(j - \frac{1}{2}, j + \frac{1}{2} \right).$$




Figure 3: Results with different discretizations. From left to right: Boundary conditions, result with only forward differences, detail of the resulting interfaces, and with the chosen upwind scheme. On the rightmost example, the resulting interfaces depend less strongly on their orientation, and there are two flip symmetries.

First, we need to decide which constraint to use in the discrete setting. We denote by

$$E - B\left(\frac{1}{n}\right) := \left\{ x \in E \mid d(x, \partial E) > \frac{1}{n} \right\},$$

Our choice is to take

$$\Omega_s^n := \bigcup_{R_{ij}^n \subset \Omega_s - B(\frac{1}{n})} R_{ij}^n$$

whereas

$$\Omega^n := [0, 1]^2 \setminus \left(\bigcup_{R_{ij}^n \subset ([0, 1]^2 \setminus \Omega) - B(\frac{1}{n})} R_{ij}^n \right),$$

such that the discrete constraints are less restrictive than the continuous ones (see Figure 4) and

$$\overline{\Omega_s^n} \Subset \Omega_s, \quad \overline{[0, 1]^2 \setminus \Omega^n} \Subset [0, 1]^2 \setminus \Omega. \quad (31)$$

We define TV^n as in [7], when the function is piecewise constant on the R_{ij}^n and $+\infty$ otherwise.

$$\text{TV}^n := \frac{1}{n^2} \sum_{i,j} |\nabla v^{ij} \vee 0|$$

with $\nabla v^{ij} \vee 0$ denotes the positive components of ∇v^{ij} , which was defined in (30), therefore picking only the ‘upwind’ variations. The norm is computed using the inner product in $\mathbb{R}^{2 \times 2}$.

We first prove the following lemma, which states that the continuous total variation may be computed with multipliers with positive components, mimicking the discrete definition.

Lemma 4. *Let $v \in \text{BV}(\mathbb{R}^d)$ and $\Omega \subset \mathbb{R}^d$ open. Then, $\text{TV}(v, \Omega) = \text{TV}^+(v, \Omega)$, where*

$$\text{TV}^+(v, \Omega) := \sup \left\{ \int_{\Omega} v \cdot \text{div } p - v \cdot \text{div } q \mid p, q \in \mathcal{C}_0^1(\Omega, \mathbb{R}^d), |p|^2 + |q|^2 \leq 1, p, q \geq 0 \right\}. \quad (32)$$

Proof. We recall that

$$\text{TV}(v, \Omega) = \sup \left\{ \int_{\Omega} v \cdot \text{div } p \mid p \in \mathcal{C}_0^1(\Omega, \mathbb{R}^d), |p| \leq 1 \right\}. \quad (33)$$




Figure 4: Discretization of the domain and constraints: The discrete grid encloses Ω , and discrete regions are only constrained if they are compactly contained in the corresponding continuous ones. Here, grey squares have free values while the black and white ones are fixed.

Let p, q be admissible in the right hand side of (32). Then we notice that $p - q$ is also admissible in (33), because p, q being componentwise positive implies

$$|p - q|^2 = |p|^2 + |q|^2 - 2p \cdot q \leq 1 - 2p \cdot q \leq 1,$$

and since $\operatorname{div}(p - q) = \operatorname{div} p - \operatorname{div} q$ we have

$$\operatorname{TV}^+(v, \Omega) \leq \operatorname{TV}(v, \Omega).$$

To prove the reverse inequality, let $\varepsilon > 0$ be arbitrary and $p_\varepsilon \in C_0^1(\Omega, \mathbb{R}^d)$ with $|p_\varepsilon| \leq 1$ such that

$$\operatorname{TV}(v, \Omega) - \int_{\Omega} \sum_{j=1}^d v_j \operatorname{div}(p_\varepsilon)_j < \varepsilon,$$

which we can write (renaming p_ε to its additive inverse, for convenience) as

$$\int_{\Omega} \left(1 - p_\varepsilon \cdot \frac{d(\nabla v)}{d|\nabla v|} \right) d|\nabla v| < \varepsilon. \quad (34)$$

Noting that $|p_\varepsilon| \leq 1$ and $\frac{d(\nabla v)}{d|\nabla v|} \leq 1$, the last inequality implies (since for $|\mu|, |\nu| \leq 1$, $|\mu - \nu|^2 \leq 2 - 2\mu \cdot \nu$) as

$$\int_{\Omega} \frac{1}{2} \left| p_\varepsilon - \frac{d(\nabla v)}{d|\nabla v|} \right|^2 d|\nabla v| < \varepsilon.$$

Notice that we may write this integral, since the function $\frac{d(\nabla v)}{d|\nabla v|}$ is a Radon-Nikodym derivative, in principle only in $L^1(\Omega, |\nabla v|)$, but its modulus is 1 for $|\nabla v|$ -almost every point [2, Corollary 1.29], so it is also in $L^2(\Omega, |\nabla v|)$. Now, by (34) and the Cauchy-Schwarz inequality we have

$$\begin{aligned} \int_{\Omega} 1 - |p_\varepsilon|^2 d|\nabla v| &= \int_{\Omega} (1 - |p_\varepsilon|)(1 + |p_\varepsilon|) d|\nabla v| \leq 2 \int_{\Omega} 1 - |p_\varepsilon| d|\nabla v| \\ &\leq 2 \int_{\Omega} \left(1 - p_\varepsilon \cdot \frac{d(\nabla v)}{d|\nabla v|} \right) d|\nabla v| < 2\varepsilon, \end{aligned} \quad (35)$$

Now we replace the components $(p_\varepsilon)_{ij}$ by $(\tilde{p}_\varepsilon)_{ij}$ which are smooth, coincide with $(p_\varepsilon)_{ij}$ out of $\{|(p_\varepsilon)_{ij}| < \sqrt{\varepsilon}\}$, that satisfy

$$|(\tilde{p}_\varepsilon)_{ij}| \leq |(p_\varepsilon)_{ij}|$$

and such that $\{(\tilde{p}_\varepsilon)_{ij} = 0\}$ is the closure of an open set: One can for example choose

$$0 < \alpha < \sqrt{\varepsilon}$$

and define a smooth nondecreasing function $\psi_\alpha : \mathbb{R} \rightarrow \mathbb{R}$ such that $\psi_\alpha(t) = t$ for $|t| \geq \alpha$, $|\psi_\alpha(t)| \leq |t|$ and $\psi_\alpha(-\alpha/2, \alpha/2) = \{0\}$ to define

$$(\tilde{p}_\varepsilon)_{ij} := \psi_\alpha \circ (p_\varepsilon)_{ij}.$$

Thus we have $|\tilde{p}_\varepsilon| \leq 1$ and $|(\tilde{p}_\varepsilon)_{ij} - (p_\varepsilon)_{ij}| \leq \sqrt{\varepsilon}$, and taking into account (34) we obtain

$$\begin{aligned} \left(\int_{\Omega} \left| \tilde{p}_\varepsilon - \frac{d\nabla v}{d|\nabla v|} \right|^2 d|\nabla v| \right)^{\frac{1}{2}} &\leq \left(\int_{\Omega} \left| p_\varepsilon - \frac{d\nabla v}{d|\nabla v|} \right|^2 d|\nabla v| \right)^{\frac{1}{2}} + \left(\int_{\Omega} |\tilde{p}_\varepsilon - p_\varepsilon|^2 d|\nabla v| \right)^{\frac{1}{2}} \\ &\leq C\sqrt{\varepsilon} (1 + |\nabla v|(\Omega)). \end{aligned} \quad (36)$$

Furthermore, using (35) and the definition of \tilde{p}_ε we obtain the estimate

$$\int_{\Omega} 1 - |\tilde{p}_\varepsilon|^2 d|\nabla v| = \int_{\Omega} 1 - |p_\varepsilon|^2 d|\nabla v| + \int_{\Omega} |p_\varepsilon|^2 - |\tilde{p}_\varepsilon|^2 d|\nabla v| \leq 2\varepsilon + 4\varepsilon|\nabla v| < C\varepsilon(1 + |\nabla v|),$$

which ensures, writing $1 - \mu : \nu = \frac{1}{2}(1 - |\mu|^2 + 1 - |\nu|^2 + |\mu - \nu|^2)$ and by (36) that

$$\left| \int_{\Omega} \left(\tilde{p}_\varepsilon : \frac{d\nabla v}{d|\nabla v|} - 1 \right) d|\nabla v| \right| \leq C\varepsilon \left(1 + (1 + |\nabla v|(\Omega))^2 \right). \quad (37)$$

Now, we notice that having fattened the level-set $\{(\tilde{p}_\varepsilon)_{ij} = 0\}$, we can write

$$(\tilde{p}_\varepsilon)_{ij} = [(\tilde{p}_\varepsilon)_{ij}]^+ - [(\tilde{p}_\varepsilon)_{ij}]^-$$

where both quantities are smooth. Writing similarly

$$\tilde{p}_\varepsilon = \tilde{p}_\varepsilon^+ - \tilde{p}_\varepsilon^-$$

with $\tilde{p}_\varepsilon^\pm$ are smooth and have only positive components, we note that $(\tilde{p}_\varepsilon^+, \tilde{p}_\varepsilon^-)$ are admissible in the right hand side of (32), so that (37) implies

$$\text{TV}^+(v, \Omega) \geq \text{TV}(v, \Omega) - C(v)\varepsilon.$$

Letting $\varepsilon \rightarrow 0$, we conclude. \square

We can now prove Gamma-convergence of the discrete problems, implying convergence of the corresponding minimizers.

Theorem 7.

$$\text{TV}^n + \chi_{C^n} \xrightarrow{\Gamma-L^1} \text{TV} + \chi_C$$

where

$$C^n := \{v = 1 \text{ on } \Omega_s^n, 0 \text{ on } [0, 1]^2 \setminus \Omega^n\}$$

and

$$C := \{v = 1 \text{ on } \Omega_s, 0 \text{ on } [0, 1]^2 \setminus \Omega\}.$$

Proof. First, we study the Γ -liminf and assume that $v_n \rightarrow v$ in L^1 . Notice that we can write $\text{TV}^n(v_n)$ as a dual formulation

$$\text{TV}^n(v_n) = \sup \{v_n \cdot \text{div}^n(p) \mid p : G \rightarrow (\mathbb{R}^2)^2\}$$

where $\text{div}^n p \in \mathbb{R}^2$ is the signed divergence corresponding to (30), and defined by

$$\begin{aligned} (\text{div}^n(p))^{i,j} := & (p_{1,+})^{i,j} - (p_{1,+})^{i-1,j} + (p_{1,-})^{i,j} - (p_{1,-})^{i+1,j} \\ & (p_{2,+})^{i,j} - (p_{2,+})^{i,j-1} + (p_{2,-})^{i,j} - (p_{2,-})^{i,j+1}. \end{aligned}$$

This is obtained easily by a (discrete) integration by parts in the expression

$$|\nabla v \vee 0| = \sup_{\substack{|p| \leq 1 \\ p_i \geq 0}} \nabla v \cdot p.$$

Now, we note that every $p : G \rightarrow (\mathbb{R}^2)^2$ can be viewed as the discretization of some smooth function $\bar{p} : [0, 1]^2 \rightarrow (\mathbb{R}^2)^2$, for example stating

$$p^{ij} = \int_{R_{ij}} \bar{p}.$$

As a result, one can write

$$\text{TV}^n(v_n) = \sup \{v_n \cdot \text{div}^n(\bar{p}) \mid \bar{p} \in C_0^1([0, 1]^2, (\mathbb{R}^2)^2), |\bar{p}| \leq 1, \bar{p} \geq 0\}.$$

It is well known that for a smooth function \bar{p} , the quantity $\text{div}^n \bar{p}$ converges to

$$\text{div} \bar{p} = \text{div}(\bar{p}_{1,1}, \bar{p}_{2,1}) + \text{div}(-\bar{p}_{1,2}, -\bar{p}_{2,2}).$$

Therefore, using Lemma 4 we get

$$\text{TV}^+(v) = \text{TV}(v) \leq \liminf \text{TV}^n(v_n).$$

For χ_{C^n} , let us first assume $\chi_C(v) = +\infty$, that is either $v \not\equiv 0$ on $[0, 1]^2 \setminus \Omega$ or $v \not\equiv 1$ on Ω_s . If the latter holds, then for ε small enough, $\Omega_s \cap (\{v > 1 + 2\varepsilon\} \cup \{v < 1 - 2\varepsilon\})$ has positive measure and thanks to the L^1 convergence of v_n ,

$$\Omega_s^n \cap (\{v_n > 1 + \varepsilon\} \cup \{v_n < 1 - \varepsilon\})$$

must have a positive measure for n big enough. That implies $\chi_{C^n}(v_n) = +\infty$ and the Γ -liminf inequality is trivially true. If $\chi_C(v) < \infty$, then $\chi_C(v) = 0$ and the inequality is also true since $C^n \subset C$.

Let now $v \in \text{BV}((0, 1)^2)$. For the Γ -limsup inequality we want to construct a sequence $v_n \rightarrow v$ such that

$$\text{TV}(v) + \chi_C(v) \geq \limsup \text{TV}^n(v_n) + \chi_{C^n}(v_n).$$

If $v \notin C$, any $v_n \rightarrow v$ gives the inequality. If $v \in C$, then we first introduce

$$v_\delta = \psi_\delta * v$$

where ψ_δ is a convolution kernel with width δ .

Then, $\text{TV}(v_\delta) \rightarrow \text{TV}(v)$ ([3, Theorem 1.3], noticing that v is constant around $\partial[0, 1]^2$) and, thanks to (31), if $\delta \leq \frac{1}{n}$, we have $\chi_{C^n}(v_\delta) = 0$.

We define $v_{\delta,n}$ by

$$(v_{\delta,n})^{ij} = \int_{R_{ij}^n} v_{\delta},$$

that satisfies $\chi_{C^n}(v_{\delta,n}) = 0$, and compute

$$\frac{|(v_{\delta,n})^{i+1,j} - (v_{\delta,n})^{i,j}|}{n} = \frac{1}{n} \left| \int_{R_{ij}^n} v_{\delta}(x + \frac{1}{n}, y, z) - v_{\delta}(x, y) \right| \geq \inf_{R_{ij}^n \cup (R_{ij}^n + (\frac{1}{n}, 0))} |\partial_x v_{\delta}|.$$

Then since $v_{\delta} \in C^1$, it is clear that the right hand side converges to $|\partial_x v_{\delta}|$. Note that in the 'upwind' gradient of a smooth function, only one term by direction can be active, then it is also true for $v_{\delta,n}$ if n is large enough and therefore $\text{TV}^n(v_{\delta,n}) \rightarrow \text{TV}(v_{\delta})$. By a diagonal argument on δ and n , we conclude. \square

5.3 Single particle results

In this section, we again restrict ourselves to the case in which there is either only one particle, or the particles are constrained to move with the same velocity.

In [9], it is shown analytically that the minimizers of TV over the set $\text{BV}_{\diamond,1}$ defined in (27) have level-sets that minimize some geometrical quantities. In particular, Theorem 4.10 there shows that there exists a minimizer of the form

$$u_0 := 1_{\Omega_1} - \lambda 1_{\Omega_-}$$

where Ω_- is the maximal Cheeger set of $\Omega \setminus \Omega_s$, and Ω_1 is a minimizer of

$$E \mapsto P(E) + \frac{P(\Omega_-)}{|\Omega_-|} |E|$$

over $E \supset \Omega_s$.

Unfortunately, determining Cheeger sets analytically is only possible in a very narrow range of sets, which makes useful the numerical computation of minimizers. We present two examples of the output of the numerical method for (28) with the constraint (29). First, we consider the "Pacman" shaped Ω_s within again a square Ω ; see Figure 5 (left). This example induces both asymmetry (left-right) and non-convexity of Ω_s which is showed in [9] to influence the geometry of the minimizer. The solution is shown in the central panel of Figure 5 and the right-hand panel shows a histogram of the solution.

The second example concerns the geometry depicted in Figure 6 (top panel), in which Ω_s denotes the two L-shaped regions in the white dumbbell-shaped domain Ω . By giving a close look, it is clear that there is a Cheeger set of $\Omega \setminus \Omega_s$ in each half of the domain, which implies the non uniqueness of the minimizer. The question is which solution the computations will converge to. Figure 6 (lower, left and right) show that different minimizers are selected numerically, in this case by using different numerical resolution.

5.4 Several particles

We now extend the numerical scheme of to optimize also over the velocities γ_i on each component Ω_s^i . The corresponding problem is again the minimization (28), but with the new constraint set

$$C^n := \left\{ v \in X \mid v = 0 \text{ on } G^n \setminus \Omega^n, v \text{ constant on } (\Omega_s^n)^i, \frac{1}{|\Omega_s^n|} \sum_{\Omega_s^n} v = 1 \right\}.$$




Figure 5: Numerical results with a single particle, illustrating the results of [9]. On the left of the two subfigures, the free part $\Omega^n \setminus \Omega_s^n$ of the computational domain G^n is in gray, and the particles Ω_s^n are white. The minimizers are on the right, where the blue colour represents the negative values and the red colour, the positive ones. The corresponding computed critical yield numbers are $Y_c = 0.0576$ and $Y_c = 0.0596$, with Ω having side length 1.

Here, $(\Omega_s^n)^i$ denotes the i -th component of the discrete domain, corresponding to Ω_s^i . The set C^n is the discrete counterpart to the set BV_\diamond used in sections 3 and 4.

We give several examples that illustrate the behavior of TV-minimizers in BV_\diamond with a disconnected Ω_s . Figure 7 shows the influence of the positions of particles with respect to each other and to the boundary, which might lump up in different configurations. Figure 8 shows two generic situations: 8(a), the flowing part is concentrated around one connected component of Ω_s whereas on 8(b), it is concentrated around the whole Ω_s .

We also give an example where uniqueness of the minimizer is not expected. In Figure 9, we consider a grid of circular particles in a square. It is easy to see analytically that any subset of the particles can be chosen as positive part of the minimizer. We present two computations at different numerical resolutions that pick two different subsets.

5.5 A random distribution of small particles

We also present two examples of random distribution of square particles in a bigger square. Figure 10 shows the same number of particles distributed in two different ways and the corresponding minimizers. This example shows that the yield number depends strongly on the geometry of the problem, not only on the ratio solid/fluid. An interesting problem would be to investigate the optimal distribution to maximize/minimize this yield number.

Acknowledgments

This research was supported by the Austrian Science Fund (FWF) through the National Research Network ‘Geometry+Simulation’ (NFN S11704). We would like to thank Ian Frigaard (UBC) for useful discussions.

References

- [1] L. Ambrosio, V. Caselles, S. Masnou, and J.-M. Morel. Connected components of sets of finite perimeter and applications to image processing. *J. Eur. Math. Soc. (JEMS)*, 3(1):39–92, 2001.




Figure 6: Boundary conditions and results computed at two different resolutions, in a situation when uniqueness of minimizers of TV in $BV_{\diamond,1}$ is not expected [17, 9]. Here $Y_c = 0.087$, where the longest side of Ω is 1.

- [2] L. Ambrosio, N. Fusco, and D. Pallara. *Functions of bounded variation and free discontinuity problems*. Oxford Mathematical Monographs. Oxford University Press, New York, 2000.
- [3] G. Anzellotti and M. Giaquinta. Existence of the displacement field for an elastoplastic body subject to Hencky’s law and von Mises yield condition. *Manuscripta Math.*, 32(1-2):101–136, 1980.
- [4] N. J. Balmforth, I. A. Frigaard, and G. Ovarlez. Yielding to stress: Recent developments in viscoplastic fluid mechanics. *Annu. Rev. Fluid Mech*, 46(1):121–146, 2014.
- [5] D. Bertsimas and J.N. Tsitsiklis. *Introduction to Linear Optimization*. Athena Scientific, 1997.
- [6] A. Chambolle, V. Duval, G. Peyré, and C. Poon. Geometric properties of solutions to the total variation denoising problem. *Inverse Prob.*, 33(1):015002, 2017.
- [7] A. Chambolle, S. E. Levine, and B. J. Lucier. An upwind finite-difference method for total variation-based image smoothing. *SIAM J. Imaging Sci.*, 4(1):277–299, 2011.
- [8] A. Chambolle and T. Pock. A first-order primal-dual algorithm for convex problems with applications to imaging. *J. Math. Imaging Vision*, 40(1):120–145, 2011.
- [9] I. A. Frigaard, J. A. Iglesias, G. Mercier, C. Pöschl, and O. Scherzer. Critical yield numbers of rigid particles settling in Bingham fluids and Cheeger sets. *SIAM J. Appl. Math.*, 77(2):638–663, 2017.
- [10] I. A. Frigaard and O. Scherzer. Uniaxial exchange flows of two Bingham fluids in a cylindrical duct. *IMA J. Appl. Math.*, 61:237–266, 1998.
- [11] I. A. Frigaard and O. Scherzer. The effects of yield stress variation in uniaxial exchange flows of two Bingham fluids in a pipe. *SIAM J. Appl. Math.*, 60:1950–1976, 2000.




Figure 7: Examples of minimizers of TV in BV_\diamond for several particles. The top row represents the boundary conditions. The computed minimizers are depicted below, where the blue colour represents the negative values and the red colour, the positive ones. The corresponding critical yield numbers are $Y_c = 0.0378$, $Y_c = 0.0383$ and $Y_c = 0.0396$ respectively, again when the longest side of Ω is 1.

- [12] R. Hassani, I. R. Ionescu, and T. Lachand-Robert. Shape optimization and supremal minimization approaches in landslides modeling. *Appl. Math. Optim.*, 52:349–364, 2005.
- [13] P. Hild, I. R. Ionescu, T. Lachand-Robert, and I. Rosca. The blocking of an inhomogeneous Bingham fluid. applications to landslides. *M2AN Math. Model. Numer. Anal.*, 36:1013–1026, 2002.
- [14] J. A. Iglesias, G. Mercier, and O. Scherzer. A note on convergence of solutions of total variation regularized linear inverse problems. Preprint arXiv:1711.06495, 2017.
- [15] I. R. Ionescu and T. Lachand-Robert. Generalized cheeger’s sets related to landslides. *Calc. Var. Partial Differential Equations*, 23:227–249, 2005.




Figure 8: Two examples of minimizing TV in BV_\diamond for several particles. Here, $Y_c = 0.0324$ (the length of a side of Ω being 1) and $Y_c = 0.0344$ (the diameter of Ω being 1). For the left result, since the magnitude of the negative values is much smaller than that of the positive ones, their color has been rescaled.




Figure 9: Numerical computation of a minimizer at two different resolutions when uniqueness is not expected. Here, since the magnitude of the negative values is much smaller than that of the positive ones, their color has been rescaled.

- [16] L. Jossic and A. Magnin. Drag and stability of objects in a yield stress fluid. *AIChE J.*, 47:2666—2672, 2001.
- [17] B. Kawohl and T. Lachand-Robert. Characterization of Cheeger sets for convex subsets of the plane. *Pacific J. Math.*, 225(1):103–118, 2006.
- [18] G. P. Leonardi and A. Pratelli. On the Cheeger sets in strips and non-convex domains. *Calc. Var. Partial Differential Equations*, 55(1):Art. 15, 28p, 2016.
- [19] E. H. Lieb and M. Loss. *Analysis*, volume 14 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2001.
- [20] P. P. Mosolov and V. P. Miasnikov. Variational methods in the theory of the fluidity of a viscous-plastic medium. *J. Appl. Math. Mech.*, 29(3):545–577, 1965.
- [21] P. P. Mosolov and V. P. Miasnikov. On stagnant flow regions of a viscous-plastic medium in pipes. *J. Appl. Math. Mech.*, 30(4):841–854, 1966.
- [22] E. Parini. An introduction to the Cheeger problem. *Surv. Math. Appl.*, 6:9–21, 2011.
- [23] A. Putz and I. A. Frigaard. Creeping flow around particles in a Bingham fluid. *J. Non-Newton. Fluid Mech.*, 165:263–280, 2010.
- [24] G. Vinay, A. Wachs, and J.-F. Agassant. Numerical simulation of non-isothermal viscoplastic waxy crude oil flows. *J. Non-Newton. Fluid Mech.*, 128(2):144 – 162, 2005.
- [25] H. F. Weinberger. Variational properties of steady fall in Stokes flow. *J. Fluid Mech.*, 52(2):321–344, 1972.




Figure 10: Two random distributions of the same number of particles in a square. On the left lies the distribution of particles and on the right the computed minimizer. Note that $Y_c^{\text{up}} = 7.89 \cdot 10^{-3}$ whereas $Y_c^{\text{low}} = 6.75 \cdot 10^{-3}$. Here, again the side length of the domain is 1 and the color of the negative values has been rescaled.